Integration of elemental and isotope data supports a Neoproterozoic Adamastor Ocean realm

F.A. Caxito^{1*}, M. Heilbron², C.M. Valeriano², H. Bruno², A. Pedrosa-Soares¹,
F.F. Alkmim³, F. Chemale⁴, L.A. Hartmann⁵, E. Dantas⁶, M.A.S. Basei⁷

doi: 10.7185/geochemlet.2106

Abstract

A robust elemental and isotopic dataset from Neoproterozoic igneous rocks discloses protracted consumption of oceanic lithosphere in the 3,000 km long orogenic system of southeastern South America. Time dependent isotopic variation trends suggest that Tonian-Cryogenian magmatic rocks formed in intra-oceanic supra-subduction settings, followed by Ediacaran magmatic arc building along Andean-type continental margins. Tectonic slices of basic-ultrabasic rocks associated with deep sea and exhalative rocks are interpreted as remnants of obducted oceanic lithosphere. Protracted closure of the oceanic realm resulted in a Himalayas-sized orogenic belt during Ediacaran-Cambrian collision, as recorded by voluminous aluminum-rich syn-collisional granites followed by post-collisional intrusions. The duration and rates of crust forming processes in island arc, continental margins and collisional settings imply that a vast Adamastor oceanic realm was consumed to form western Gondwana.

Received 23 September 2020 | Accepted 6 January 2021 | Published 11 February 2021

Introduction

The Cretaceous breakup of Gondwana in the South Atlantic region produced an asymmetric division of the Neoproterozoic Brasiliano/Pan-African Orogen (Fig. 1a,b) (Cordani *et al.*, 2003). A large part of the orogen remained in South America as the 3,000 km long and 100–500 km wide Mantiqueira Province (Almeida *et al.*, 1981) (Fig. 1a).

Early postulations of tectonic evolution of the Mantiqueira Province emphasised intra-continental models (*e.g.*, Torquato and Cordani, 1981; Porada *et al.*, 1989), an interpretation that received renewed attention in recent years (Cavalcante *et al.*, 2019; Meira *et al.*, 2019; Fossen *et al.*, 2020; Konopásek *et al.*, 2020). However, a wealth of field, geochemical and geochronological data produced in recent decades progressively prompted interpretations involving distinct stages of typical Wilson cycle processes involved in the closure of the Adamastor Ocean (originally defined by Hartnady *et al.*, 1985). Besides reworked Archean-Palaeoproterozoic basement, Neoproterozoic continental rift and passive margin successions, ocean floor assemblages and syn-orogenic basins were described. Those are intruded by large volumes of Cryogenian to Cambrian plutonic rocks typical of pre-collisional, collisional, and post-collisional stages (Fig. 1a), with temporal and spatial

ordering leading to the present unraveling of a simple and reliable interpretation of a complete rift-drift-subduction-collision plate tectonic cycle. The archetypical examples of intracontinental orogens (Raimondo *et al.*, 2014), as proposed in the last century and recently revived, lack many of the tectonic components recognised in the last decades in the Mantiqueira Province.

Here we provide a new interpretation for a comprehensive database of 1,583 Hf-in-zircon (839 from Neoproterozoic, 744 from Archean-Palaeoproterozoic basement), 358 Sm-Nd isotope (221 and 137) and 824 bulk rock elemental geochemistry determinations of Neoproterozoic plutonic and volcanic rocks of the Mantiqueira Province (Supplementary Information), and demonstrate that these rocks are a testimony to an oceanic realm that was consumed to generate a Himalayas-sized orogen.

Tonian-Cryogenian assemblages with Volcanic Arc Signatures

Tonian-Cryogenian tonalite-granodiorite orthogneisses with dioritic to mafic enclaves represent an expanded series of calc-alkaline, magnesian, metaluminous, I-type magmas along tholeiitic basalt (Fig. 1c-f). The 835–860 Ma Serra da Prata Complex of the central province has whole rock ϵNd_0 from +6.4 to +0.9

1. Universidade Federal de Minas Gerais, CPMTIC-IGC, Belo Horizonte, 31270-901 MG, Brazil
2. Universidade do Estado do Rio de Janeiro, Faculdade de Geologia, Rio de Janeiro, 20559-900 RJ, Brazil
3. Universidade Federal de Ouro Preto, Departamento de Geologia, Ouro Preto, 35400-000 MG, Brazil
4. Universidade do Vale do Rio dos Sinos, Faculdade de Geologia, São Leopoldo, 93022-750 RS, Brazil
5. Universidade Federal do Rio Grande do Sul, Instituto de Geociências, Porto Alegre, 91501-970 RS, Brazil
6. Universidade de Brasília, Laboratório de Geocronologia, Brasília, 70910-900 DF, Brazil
7. Universidade de São Paulo, CPGeo-IGC, São Paulo, 05508-080 SP, Brazil

* Corresponding author (email: caxito@ufmg.br)

Figure 1 (a) Geological map of the Mantiqueira Province of southeastern South America in the context of (b) West Gondwana, with (c-f) elemental and (g-j) isotopic data from basic to felsic plutonic rocks and mafic-ultramafic bodies (see [Supplementary Information](#) for sources and references).

with T_{DM} Nd of 860–1,200 Ma, $^{87}\text{Sr}/^{86}\text{Sr}_i < 0.7035$, zircon ϵHf_i from +14 to +10, and T_{DM} Hf of 840–1,010 Ma (Peixoto *et al.*, 2017; Heilbron *et al.*, 2020; Santiago *et al.*, 2020). The Rio

Negro Complex comprises intermediate to felsic plutonic rocks (620–790 Ma) with ϵNd_i from +5 to –3 and $^{87}\text{Sr}/^{86}\text{Sr} < 0.705$, as well as high K calc-alkaline to shoshonitic felsic rocks

(605–620 Ma) with $\epsilon\text{Nd}_{(t)} = -3$ to -14 and $^{87}\text{Sr}/^{86}\text{Sr} = 0.7050\text{--}0.7100$ (Tupinambá *et al.*, 2012).

The São Gabriel terrane in the southern region (Fig. 1a) yielded U–Pb zircon ages of 800–860 Ma and 680–770 Ma (Babinski *et al.*, 1996; Saalman *et al.*, 2005). Sm–Nd T_{DM} are at 800–1,000 Ma with $\epsilon\text{Nd}_{(t)}$ on the depleted mantle curve (up to $+8$, Babinski *et al.*, 1996; Saalman *et al.*, 2005) and zircon $\epsilon\text{Hf}_{(t)}$ from $+14$ to $+8$ (Cerva-Alves *et al.*, 2020).

Ediacaran Assemblages with Volcanic Arc Signatures

Widespread Ediacaran (mainly *ca.* 580–630 Ma) magmatic rocks represent an expanded series of medium to high K calc-alkaline, magnesian, metaluminous, I-type tonalites to granodiorites rich in dioritic to mafic enclaves, with minor gabbro (Fig. 1a) (Tedeschi *et al.*, 2016; Basei *et al.*, 2018; Corrales *et al.*, 2020). The Rio Doce plutonic and volcanic rocks show $\epsilon\text{Nd}_{(t)}$ of -2.9 to -13.6 with $T_{\text{DM}}\text{Nd}$ of 1,190–2,030 Ma, $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.7059–0.7165 and zircon $\epsilon\text{Hf}_{(t)}$ of $+2.3$ to -11.7 (Tedeschi *et al.*, 2016; Degler *et al.*, 2017; Novo *et al.*, 2018; Araújo *et al.*, 2020; Corrales *et al.*, 2020). The Pelotas-Florianópolis-Aiguá batholith forms a multi-intrusion geological structure consisting of granite, gabbro and diorite with $^{87}\text{Sr}/^{86}\text{Sr}_i$ ratios of *ca.* 0.712, ϵNd of -3.6 to -22.2 and $T_{\text{DM}}\text{Nd}$ of 1,200–2,400 Ma (Babinski *et al.*, 1997; Koester *et al.*, 2016).

Neoproterozoic Basic-Ultrabasic Assemblages

Accretionary mélanges of the northern province (Fig. 1a) include MORB chemistry metabasalt, banded metadolerite, metagabbro and meta-ultramafic rocks, with $\epsilon\text{Nd}_{(t)}$ up to $+6.3$ (Pedrosa-Soares *et al.*, 1998; Amaral *et al.*, 2020). Rootless plagiogranite veins hosted by a banded metadolerite are dated at 645 ± 10 Ma (Amaral *et al.*, 2020).

Mafic-ultramafic rock associations in the central sector include dunite cumulates, MORB- and IAT-like gabbro, metabasic rocks with sheeted dikes, pillow lavas and chert, emplaced at *ca.* 630 Ma (Tassinari *et al.*, 2001; Passarelli *et al.*, 2018).

The 715–920 Ma MORB chemistry mafic-ultramafic assemblages in the southern sector have zircon $\epsilon\text{Hf}_{(t)}$ up to $+15$ and mantle-like trace element signatures (Arena *et al.*, 2017, 2018; Hartmann *et al.*, 2019). Dravite in altered oceanic crust has typical ocean floor $\delta^{11}\text{B}$ up to $+1.8$ (Hartmann *et al.*, 2019). The southernmost ophiolites show whole rock $\epsilon\text{Nd}_{(t)}$ up to $+8.5$ (Peel *et al.*, 2018; Ramos *et al.*, 2020) (Fig. 1a).

Discussion

The time dependent variation of isotopic trends (Fig. 1g–j) is interpreted as a shift from Tonian–Cryogenian juvenile settings fingerprinting intra-oceanic arcs to Ediacaran settings with magmas formed by mixing of mantle wedge melts and anatectic melts from both the Tonian–Cryogenian rocks and the Archean–Palaeoproterozoic basement. The resulting mixed melts intruded active continental margin settings similar to Andean volcanic arcs. The protracted consumption of oceanic lithosphere is required, and this is attested by the associated ophiolites along the entire length of the Mantiqueira Province.

Recently revisited models of intracontinental orogeny focused on a “space problem” for the development of an Adamastor oceanic realm based on estimates of maximum

oceanic width using modern spreading and subduction rates (Fossen *et al.*, 2020; Konopásek *et al.*, 2020). However, those estimates are contingent and cannot be used as hard proof against mobilistic models, because: 1) the ocean was probably fragmented in smaller sub-domains such as in the present day western Pacific and also connected to a much large Neoproterozoic oceanic system (Cordani *et al.*, 2003) (Fig. 1b); 2) the Neoproterozoic upper mantle was warmer, leading to distinct spreading and subduction rates and dynamics (Brown, 2014); 3) intense wrench tectonics with thousands-of-km long shear zones obliterated former low angle surfaces and displaced allochthonous units far from their original position; 4) age, size and time span of an ocean have little significance in determining whether subduction is feasible or not (Hall, 2019), while lithospheric weaknesses such as magma-rich rift systems present in the orogen precursor basins (Tack *et al.*, 2001; Basei *et al.*, 2008) might play a major role in subduction initiation (Stern and Gerya, 2018); 5) those calculations use a palinspastic incorrect pre-drift reconstruction, ignoring at least *ca.* 280 km of restored continental crust now stretched in the Brazilian and African passive margins (Aslanian *et al.*, 2009); 6) the termination of the orogen in a gulf partially enclosed by continental crust implies much lower convergence rates; and 7) non-Euclidean geometry implies the diachronous along strike opening and closure of oceanic basins. Thus, calculations of former oceanic width are interesting hypothetical exercises, but the field, petrographic, isotopic, elemental and geochronological data sets cannot be considered as subordinated to it.

The striking coherence of evidence from various geological disciplines (detailed field, petrographic and structural studies, bulk rock chemistry, isotope geochemistry, quantitative geothermobarometry and petrology) support a solid interpretation for the tectonic units and their arrangement in time and space to be explained better through the development and consumption of an oceanic realm in the Mantiqueira Province. The evidence provided here suggests that the main ocean was located to the west of the arc rocks, where recently paired HP-HT metamorphic belts were described (Ricardo *et al.*, 2020). The original proposition of Hartnady *et al.* (1985) would correspond to the Marmora back-arc basin in the African counterparts, east of the Pelotas arc (Frimmel *et al.*, 2011; Basei *et al.*, 2018).

Acknowledgements

We dedicate this article to the memory of Koji Kawashita, a pioneering, devoted scientist who dedicated his career to the implementation of many isotope geology laboratories in Brazil. The authors acknowledge the Brazilian Scientific and Technological Development Council (CNPq) for their research grants. Comments by Craig Storey and Hartwig Frimmel are acknowledged. This is a contribution to Project MOBILE (geolifemobile.com) supported by Instituto Serrapilheira (Serra-1912-31510).

Editor: Horst R. Marschall

Additional Information

Supplementary Information accompanies this letter at <https://www.geochemicalperspectivesletters.org/article2106>.

© 2021 The Authors. This work is distributed under the Creative Commons Attribution Non-Commercial No-Derivatives 4.0

License, which permits unrestricted distribution provided the

original author and source are credited. The material may not be adapted (remixed, transformed or built upon) or used for commercial purposes without written permission from the author. Additional information is available at <https://www.geochemicalperspectivesletters.org/copyright-and-permissions>.

Cite this letter as: Caxito, F.A., Heilbron, M., Valeriano, C.M., Bruno, H., Pedrosa-Soares, A., Alkmim, F.F., Chemale, F., Hartmann, L.A., Dantas, E., Basei, M.A.S. (2021) Integration of elemental and isotope data supports a Neoproterozoic Adamastor Ocean realm. *Geochem. Persp. Let.* 17, 6–10.

References

- ALMEIDA, F.F.M., HASUI, Y., BRITO NEVES, B.B., FUCK, R.A. (1981) Brazilian structural provinces: an introduction. *Earth-Science Reviews* 17, 1–29.
- AMARAL, L., CAXITO, F.A., PEDROSA-SOARES, A.C., QUEIROGA, G., BABINSKI, M., TRINDADE, R., LANA, C., CHEMALE, F. (2020) The Ribeirão da Folha ophiolite-bearing accretionary wedge (Araçuai orogen, SE Brazil): new data for Cryogenian plagiogranite and metasedimentary rocks. *Precambrian Research* 336, 105522.
- ARAÚJO, C., PEDROSA-SOARES, A., LANA, C., TEDESCHI, M., DUSSIN, I. (2020) Primeiro registro de magmatismo juvenil no arco magmático Rio Doce, Orógeno Araçuai meridional. In: XVII Simpósio Nacional de Estudos Tectônicos (SNET), SBG, Bento Gonçalves, Annals, 105.
- ARENA, K.R., HARTMANN, L.A., LANA, C. (2017) Tonian emplacement of ophiolites in the southern Brasiliano Orogen delimited by U-Pb-Hf isotopes of zircon from metasomatites. *Gondwana Research* 49, 296–332.
- ARENA, K.R., HARTMANN, L.A., LANA, C. (2018) U-Pb-Hf isotopes and trace elements of metasomatic zircon delimit the evolution of neoproterozoic Capané ophiolite in the southern Brasiliano Orogen. *International Geology Review* 60, 911–928.
- ASLANIAN, D., MOULIN, M., OLIVET, J. L., UNTERNEHR, P., MATIAS, L., BACHE, F., RABINEAU, M., NOUZÉ, H., KLINGELHOEFER, F., CONTRUCCI, I., LABAILS, C. (2009) Brazilian and African passive margins of the Central Segment of the South Atlantic Ocean: Kinematic constraints. *Tectonophysics* 468, 98–112.
- BABINSKI, M., CHEMALE, F., HARTMANN, L.A., VAN SCHMUS, W.R., SILVA, L.C. (1996) Juvenile accretion at 750–700 Ma in southern Brazil. *Geology* 24, 439–442.
- BABINSKI, M., CHEMALE, F., VAN SCHMUS, W.R., HARTMANN, L.A., SILVA, L.C. (1997) U-Pb and Sm-Nd geochronology of the neoproterozoic granitic-gneissic Dom Feliciano Belt, southern Brazil. *Journal of South American Earth Sciences* 10, 263–274.
- BASEI, M.A.S., FRIMMEL, H.E., NUTMAN, A.P., PRECIOZZI, F. (2008) West Gondwana amalgamation based on detrital zircon ages from Neoproterozoic Ribeira and Dom Feliciano belts of South America and comparison with coeval sequences from SW Africa. *Geological Society, London, Special Publications* 294, 239–256.
- BASEI, M.A.S., FRIMMEL, H.E., CAMPOS NETO, M.C., GANADE DE ARAÚJO, C.E., CASTRO, N.A., PASSARELLI, C.R. (2018) The tectonic history of the southern Adamastor Ocean based on a correlation of the Kaoko and Dom Feliciano belts. In: SIEGISMUND S., BASEI M., OYHANTCABAL P., ORIOLO S. (Eds.) *Geology of Southwest Gondwana. Regional Geology Reviews*. Springer, Cham., 63–85.
- BROWN, M. (2014) The contribution of metamorphic petrology to understanding lithosphere evolution and geodynamics. *Geoscience Frontiers* 5, 553–569.
- CAVALCANTE, C., FOSSEN, H., ALMEIDA, R.P., HOLLANDA, M.H.B., EGYDIO-SILVA, M. (2019) Reviewing the puzzling intracontinental termination of the Araçuai-West Congo orogenic belt and its implications for orogenic development. *Precambrian Research* 322, 85–98.
- CERVA-ALVES, T., HARTMANN, L.A., REMUS, M.V.D., LANA, C. (2020) Integrated ophiolite and arc evolution, southern Brasiliano Orogen. *Precambrian Research* 341, 105648.
- CORDANI, U.G., BRITO NEVES, B.B., D'AGRELLA-FILHO, M.S. (2003) From Rodinia to Gondwana: a review of the available evidence from South America. *Gondwana Research* 6, 275–283.
- CORRALES, F.F., DUSSIN, I.A., HEILBRON, M., BRUNO, H., BERSAN, S., VALERIANO, C.M., PEDROSA-SOARES, A.C., TEDESCHI, M. (2020) Coeval high Ba-Sr arc-related and OIB Neoproterozoic rocks linking pre-collisional magmatism of the Ribeira and Araçuai orogenic belts, SE-Brazil. *Precambrian Research* 337, 105476.
- DEGLER, R., PEDROSA-SOARES, A., DUSSIN, I., QUEIROGA, G., SCHULZ, B. (2017) Contrasting provenance and timing of metamorphism from paragneisses of the Araçuai-Ribeira orogenic system, Brazil: Hints for Western Gondwana assembly. *Gondwana Research* 51, 30–50.
- FOSSEN, H., CAVALCANTE, C., KONOPÁSEK, J., MEIRA, V.T., ALMEIDA, R.P., HOLLANDA, M.H.B., TROMPETTE, R. (2020) A critical discussion of the subduction-collision model for the Neoproterozoic Araçuai-West Congo orogen. *Precambrian Research* 343, 105715.
- FRIMMEL, H.E., BASEI, M.S., GAUCHER, C. (2011) Neoproterozoic geodynamic evolution of SW-Gondwana: a southern African perspective. *International Journal of Earth Sciences* 100, 323–354.
- HALL, R. (2019) The subduction initiation stage of the Wilson cycle. *Geological Society, London, Special Publications* 470, 415–437.
- HARTMANN, L.A., WERLE, M., MICHELIN, C.R.L., LANA, C., QUEIROGA, G., CASTRO, M.P., ARENA, K.R. (2019) Proto-Adamastor ocean crust (920 Ma) described in Brasiliano Orogen from coetaneous zircon and tourmaline. *Geoscience Frontiers* 10, 1623–1633.
- HARTNADY, C., JOUBERT, P., STOWE, C. (1985) Proterozoic crustal evolution in southwestern Africa. *Episodes* 8, 236–244.
- HEILBRON, M., VALERIANO, C.M., PEIXOTO, C., TUPINAMBÁ, M., NEUBAUER, F., DUSSIN, I., CORRALES, F., BRUNO, H., LOBATO, M., ALMEIDA, J.C.H., SILVA, L.G.E. (2020) Neoproterozoic magmatic arc systems of the central Ribeira belt, SE-Brazil, in the context of the West-Gondwana pre-collisional history: A review. *Journal of South American Earth Sciences*, 102710.
- KOESTER, E., PORCHER, C. C., PIMENTEL, M. M., FERNANDES, L.A.D., VIGNOLLE-LARGE, M.L., OLIVEIRA, L.D., RAMOS, R.C. (2016) Further evidence of 777 Ma subduction-related continental arc magmatism in Eastern Dom Feliciano Belt, southern Brazil: the Chácara das Pedras Orthogneiss. *Journal of South American Earth Sciences* 68, 155–166.
- KONOPÁSEK, J., CAVALCANTE, C., FOSSEN, H., JANOUŠEK, V. (2020) Adamastor—an ocean that never existed?. *Earth-Science Reviews* 205, 103201.
- MEIRA, V.T., GARCIA-CASCO, A., HYPPOLITO, T., JULIANI, C., SCHORSCHER, J.H.D. (2019) Tectono-Metamorphic Evolution of the Central Ribeira Belt, Brazil: A Case of Late Neoproterozoic Intracontinental Orogeny and Flow of Partially Molten Deep Crust During the Assembly of West Gondwana. *Tectonics* 38, 3182–3209.
- NOVO, T.A., PEDROSA-SOARES, A., VIEIRA, V.S., DUSSIN, I., SILVA, L.C. (2018) The Rio Doce Group revisited: an Ediacaran arc-related volcano-sedimentary basin, Araçuai orogen (SE Brazil). *Journal of South American Earth Sciences* 85, 345–361.
- PASSARELLI, C.R., BASEI, M.A.S., SIGA JR., O., HARARA, O.M.M. (2018) The Luis Alves and Curitiba terranes: continental fragments in the Adamastor Ocean. In: SIEGISMUND S., BASEI M., OYHANTCABAL P., ORIOLO S. (Eds) *Geology of Southwest Gondwana. Regional Geology Reviews*. Springer, Cham., 189–215.
- PEDROSA-SOARES, A., VIDAL, P., LEONARDO, O.H., BRITO NEVES, B.B. (1998) Neoproterozoic oceanic remnants in eastern Brazil: further evidence and refutation of an exclusively ensialic evolution for the Araçuai–West Congo orogen. *Geology* 26, 519–522.
- PEEL, E., SANCHEZ-BEITUCCI, L., BASEI, M.A.S. (2018) Geology and geochronology of Paso del Dragón Complex (northeastern Uruguay): Implications on the evolution of the Dom Feliciano Belt (Western Gondwana). *Journal of South American Earth Sciences* 85, 250–262.
- PEIXOTO, C.A., HEILBRON, M., RAGATKY, D., ARMSTRONG, R., DANTAS, E., VALERIANO, C.M., SIMONETTI, A. (2017) Tectonic evolution of the Juvenile Tonian Serra da Prata magmatic arc in the Ribeira belt, SE Brazil: Implications for early west Gondwana amalgamation. *Precambrian Research* 302, 221–254.
- PORADA, H. (1989) Pan-African rifting and orogenesis in southern to equatorial Africa and eastern Brazil. *Precambrian Research* 44, 103–136.
- RAMOS, R.C., KOESTER, E., VIEIRA, D.T. (2020) Sm–Nd systematics of metatramafic rocks from the Arroio Grande Ophiolite (Brazil): Insights on the evolution of the South Adamastor paleo-ocean. *Geoscience Frontiers*, doi: 10.1016/j.gsf.2020.02.013.
- RAIMONDO, T., HAND, M., COLLINS, W.J. (2014) Compressional intracontinental orogens: Ancient and modern perspectives. *Earth-Science Reviews* 130, 128–153.
- RICARDO, B.S., FALEIROS, F.M., MORAES, R., SIGA JR., O., CAMPANHA, G.A. (2020) Tectonic implications of juxtaposed high- and low-pressure metamorphic field gradient rocks in the Turvo-Cajati Formation, Curitiba Terrane, Ribeira Belt, Brazil. *Precambrian Research*, 105766.
- SAALMANN, K., HARTMANN, L.A., REMUS, M.V.D., KOESTER, E., CONCEIÇÃO, R.V. (2005) Sm–Nd isotope geochemistry of metamorphic volcano-sedimentary successions in the São Gabriel Block, southernmost Brazil: evidence for the existence of juvenile Neoproterozoic oceanic crust to the east of the Rio de la Plata craton. *Precambrian Research* 136, 159–175.

- SANTIAGO, R., CAXITO, F.A., PEDROSA-SOARES, A., NEVES, M., DANTAS, E. (2020) Tonian island arc remnants in the northern Ribeira orogeny of western Gondwana: The Caxixe batholith (Espírito Santo state, SE Brazil). *Precambrian Research* 351, 105944.
- STERN, R.J., GERYA, T. (2018) Subduction initiation in nature and models: A review. *Tectonophysics* 746, 173–198.
- TACK, L., WINGATE, M.T.D., LIÉGEOIS, J.P., FERNANDEZ-ALONSO, M., DEBLOND, A. (2001) Early Neoproterozoic magmatism (1000–910 Ma) of the Zadinian and Mayumbian Groups (Bas-Congo): onset of Rodinia rifting at the western edge of the Congo craton. *Precambrian Research* 110, 277–306.
- TASSINARI, C.C., MUNHÁ, J.M., RIBEIRO, A., CORREIA, C.T. (2001) Neoproterozoic oceans in the Ribeira Belt (southeastern Brazil): The Pirapora do Bom Jesus ophiolitic complex. *Episodes*, 24, 245–251.
- TEDESCHI, M., PEDROSA-SOARES, A., DUSSIN, I., TASSINARI, C., SILVA, L.C., GONÇALVES, L.E., ALKIM, L., LANA, C., FIGUEIREDO, C., DANTAS E., MEDEIROS, S., CAMPOS, C., CORRALES, F., HEILBRON, M. (2016) The Ediacaran Rio Doce magmatic arc revisited (Araçuaí-Ribeira orogenic system, SE Brazil). *Journal of South American Earth Sciences* 68, 167–186.
- TORQUATO, J.R., CORDANI, U.G. (1981) Brazil-Africa geological links. *Earth-Science Reviews* 17, 155–176.
- TUPINAMBÁ, M., HEILBRON, M., VALERIANO, C., PORTO JR., R., DIOS, F.B., MACHADO, N., SILVA, L.G.E., ALMEIDA, J.C.H. (2012) Juvenile contribution of the Neoproterozoic Rio Negro magmatic arc (Ribeira Belt, Brazil): implications for western Gondwana amalgamation. *Gondwana Research* 21, 422–438.