


Oxygen limitation can trigger the production of branched GDGTs in culture

T.A. Halamka^{1*}, J.M. McFarlin¹, A.D. Younkin¹, J. Depoy¹, N. Dildar¹, S.H. Kopf¹



doi: 10.7185/geochemlet.2132

Abstract



Branched glycerol dialkyl glycerol tetraethers (brGDGTs) are ubiquitous and well-preserved sedimentary biomarkers. These compounds serve as important palaeoenvironmental indicators due to strong empirical correlations between brGDGT distributions and temperature and pH in modern environments. However, the mechanistic link between temperature, pH, and brGDGT production has been difficult to study due to uncertainty about the organisms and physiological conditions required for the synthesis of brGDGTs in nature. Here, we report that in the Acidobacterium *Edaphobacter aggregans*, severe oxygen limitation (1% O₂) leads to significant brGDGT production (2.9% of membrane lipids) with three structural varieties including an uncharacterized isomer of brGDGT Ic. Low O₂ as a

physiological trigger for brGDGT production may help explain why brGDGT producers have been so difficult to identify, and provides a new path towards uncovering the genetic basis and biological function of brGDGTs. If the oxygen effects observed here apply more broadly, the empirical calibrations for brGDGT-based temperature and pH reconstructions may currently be missing the effects of oxygen as a relevant and possibly dominant control in the environmental distributions of brGDGTs.

Received 4 May 2021 | Accepted 14 September 2021 | Published 3 November 2021

Introduction

Methodological advances in sample preparation and analysis over the past decade have highlighted the vast global distribution of brGDGTs across terrestrial, aquatic, hydrothermal, and sedimentary systems (Lincoln *et al.*, 2013; De Jonge *et al.*, 2014; Weber *et al.*, 2018; Wang *et al.*, 2019). The relative abundances of structurally unique brGDGTs have been explored in many of these settings in efforts to calibrate brGDGTs as a palaeoenvironmental proxy by establishing empirical correlations with temperature and pH, such as the Methylation index of Branched Tetraethers (MBT) and the Cyclization index of Branched Tetraethers (CBT), respectively (Weijers *et al.*, 2007; Peterse *et al.*, 2012; Naafs *et al.*, 2017). Prior work has demonstrated that many Acidobacteria, a diverse and widespread phylum of soil bacteria, synthesise the potential brGDGT precursor iso-diabolic acid (13,16-dimethyl octacosanedioic acid) in large quantities (Sinninghe Damsté *et al.*, 2018), and that some members of subdivision 1 (SD 1) Acidobacteria, including *Edaphobacter aggregans*, produce trace amounts of at least one brGDGT (Sinninghe Damsté *et al.*, 2011). However, the lack of cultured organisms that consistently produce branched tetraethers raises the question of how these compounds are so structurally diverse and abundant in nature yet so elusive in the laboratory.

Here we investigated the effects of molecular oxygen availability (O₂) on brGDGT production to test the hypothesis that brGDGT production may require a specific environmental

constraint. Many Acidobacteria including *E. aggregans* harbour high affinity terminal oxidases in their genomes (Eichorst *et al.*, 2018). These types of oxidases often have half-saturation constants at low nM concentrations of O₂, likely enabling survival and growth in micro-aerobic habitats (Pitcher and Watmough, 2004). Such low O₂ availability is common in many soil, peat, and sedimentary environments and prior work on brGDGT distribution and production across oxygen gradients suggests that some environmental source organisms may preferentially grow at oxic/anoxic transitions (Liu *et al.*, 2014; Weber *et al.*, 2018; Martínez-Sosa and Tierney, 2019). To study the effects of O₂ limitation, we examined the tetraether and fatty acid membrane composition of *E. aggregans* grown in a simplified yeast extract medium under O₂ conditions ranging from fully aerated (21 % O₂) to severely O₂ limited (1 % O₂). An excess additional carbon source (sucrose) was either added or omitted to control for the potential effects of growth rate.

Results and Discussion

The effects of O₂ limitation. *E. aggregans* produced similar fatty acids under all growth conditions with the sum of just four fatty acids (iso-diabolic acid, iso-C15:0, C16:0, and C16:1) constituting over 90 % of the fatty acid fraction (Fig. 1, Table S-1), consistent with previous observations in a different growth medium (91 %; Sinninghe Damsté *et al.*, 2011). Although the relative abundance of individual fatty acids differed significantly between culture conditions, the hypothesised brGDGT precursor

1. Department of Geological Sciences, University of Colorado, Boulder, Colorado 80309, USA

* Corresponding author (email: toby.halamka@colorado.edu)






Figure 1 Growth rate and lipid membrane profiles of *E. aggregans* grown in liquid culture at 21 % O₂ (= air, left) and 1 % O₂ (right) with and without carbon excess. Data points represent averages and standard deviations of biological triplicates. **(a)** Growth rates with medium containing 1 g/L Yeast Extract (YE, circle) and 1 g/L YE + 30 g/L sucrose (triangle). **(b)** Tetraethers [% of all detected core lipids]: brGTGT Ia (green), brGDGT Ic isomer (blue), brGDGT Ia (orange). No tetraethers were detected at 21 % O₂. **(c)** Most abundant FAMES (Fatty Acid Methyl Esters, % of all detected core lipids): iso-C15:0 (light yellow), C16:0 (red), iso-diabolic acid (purple).

iso-diabolic acid was always the dominant membrane core lipid ranging in relative abundance from 40 ± 3 % to 84 ± 2 % with a systematic decrease both at lower O₂ and without sucrose. In liquid culture, the organism produced no detectable tetraethers at 21 % and 5 % O₂, however, at 1 % O₂ *E. aggregans* produced significant quantities (up to 2.9 ± 0.7 %) of up to three different tetraethers (brGDGT Ia, brGTGT Ia, and an uncharacterised isomer of brGDGT Ic; see Fig. 2 for structures).

Growth rates slowed significantly at lower O₂ when grown with excess sucrose (from 0.89 ± 0.02 /day at 21 % O₂ to 0.50 ± 0.10 /day at 1 % O₂) and remained unchanged when grown without sucrose (0.57 ± 0.03 /day at 21 % O₂ to 0.56 ± 0.06 /day at 1 % O₂) but tetraether production in liquid culture was limited to 1 % O₂ conditions independent of growth rate (Fig. 1, Table S-2). Temperature and pH were identical for all culturing conditions, suggesting that O₂ limitation is the primary environmental trigger for brGDGT production in *E. aggregans*, with a threshold for biosynthesis in liquid culture between 5 % and 1 % O₂. However, *E. aggregans*, named for its tendency to aggregate, displays significant cellular clumping in liquid culture. Cell aggregates form natural O₂ gradients (Wessel *et al.*, 2014) and may trigger brGDGT biosynthesis by reducing O₂ availability to individual cells deeper within the aggregates. Consequently, the actual biochemical O₂ threshold for brGDGT production may be significantly lower.

Although the physiological role of brGDGTs is not yet understood, we speculate that brGDGT production in *E. aggregans* could be part of an energy stress response imposed by the lack of sufficient O₂ for energy production. Such terminal electron acceptor limitation could place increased importance on the

role of lipids in restricting membrane permeability. In Archaea, which produce isoprenoidal GDGTs (isoGDGTs), tetraether lipids are often inferred to play a similar role in combatting energy stress by decreasing the cellular membrane's proton permeability and thus decreasing maintenance energy costs (Qin *et al.*, 2015). Recent work also suggests that the synthesis of ether bonds under oxygen limiting conditions is an energetic adaptation to reducing environments (Boyer *et al.*, 2020). The structural attributes of the tetraether lipids that *E. aggregans* produces that may play a role in decreased membrane permeability include their membrane spanning nature and ether bonds. Specifically, membrane spanning lipids increase membrane rigidity and stability and ether-bound lipids have been found to decrease membrane permeability relative to ester-bound lipids (Van de Vossenberg *et al.*, 1998).

Tetraether structures. The tetraethers of *E. aggregans* were identified based on their retention times and fragmentation patterns as brGDGT Ia, brGTGT Ia (Glycerol Trialkyl Glycerol Tetraether), and a structural isomer of brGDGT Ic (Fig. 2). Retention time offsets between the newly described tetraethers (brGDGT Ic isomer and brGTGT Ia) and previously described structurally similar tetraethers (brGDGT Ic and C46 GTGT) are shown in Figure 2b. These data confirm and quantify for the first time the previously suspected production of brGDGT Ia in *E. aggregans* (Sinninghe Damsté *et al.*, 2011), and reveal the capability of this species to produce other forms of tetraethers including a potentially cyclised structure. However, the brGDGT Ic isomer we observe is asymmetric with two unsaturation equivalents on the same alkyl chain. BrGDGT Ic commonly found in environmental samples contains only one unsaturation (a pentacyclic ring) on each alkyl chain instead. While the unsaturation equivalents could stem from cyclisation pointing to an isomer of brGDGT Ic with a bicyclic alkyl chain, other structural variations such as double bonds and cyclohexyl rings could produce the observed mass spectrum (see Fig. 2 for two potential structures). While hydrogenation of the tetraethers did not reduce the brGDGT Ic isomer to brGDGT Ia, suggesting cyclisation instead of double bonds, tetraethers with cyclopentyl rings usually elute later under normal phase chromatography (Liu *et al.*, 2016), not earlier as observed here. Future analyses by NMR or ether cleavage and GC-MS with a purified concentrated fraction of this trace membrane component will help determine the exact structure of this new brGDGT Ic isomer. If *E. aggregans* indeed produces a cyclised brGDGT it must contain a cyclisation pathway different from the archaea, as the previously described archaeal grsAB tetraether cyclisation genes (Zeng *et al.*, 2019) do not have homologues in the *E. aggregans* genome. The third tetraether found in *E. aggregans*, brGTGT Ia, is likewise rarely identified in environmental samples but other tri-alkyl tetraethers have been previously observed. Although the exact biosynthetic pathway for branched tetraethers remains unresolved, we note the correlation between the occurrence of brGTGT Ia and elevated quantities of iso-C15.

Environmental implications. The clumping phenotype of *E. aggregans* highlights the potential importance of micro-scale spatial O₂ gradients in brGDGT production and is supported by *E. aggregans* growth on solid medium; unlike in liquid cultures, we observed measurable quantities of both brGDGT Ia and the brGDGT Ic isomer in fully oxygenated (21 % O₂) plate growth experiments (Table S-1). Aerobic plate growth and colony formation often produce micro-aerophilic environments within colonies, thus a significant portion of the plate culture likely experienced severe O₂ limitation thereby triggering brGDGT production. This mode of growth on a solid substrate, rather than in liquid culture, is much more representative of the lifestyle of soil microorganisms in their natural environment (Kolter and




Figure 2 MS2 fragmentation and peak retention time patterns of tetraether lipids identified in *E. aggregans* using normal phase liquid chromatography. (a) MS2 fragmentation of brGTGT Ia (green), brGDGT Ic isomer (blue), and brGDGT Ia (orange). Fragment masses and formulae shown in black. Losses from the molecular ion ($M+H-x$) shown in corresponding compound colour. Structures of three key identifying fragments included. (b) Peak retention times of the tetraethers identified in *E. aggregans* in comparison to structurally similar tetraethers from a soil standard and the C46 GTGT analytical standard (grey). Peak intensities are normalised.

Greenberg, 2006) and speaks to the potential importance of O_2 limitation in heterogenous natural settings.

To test if all Acidobacteria exhibit similar suboxia-induced brGDGT production, we conducted follow up experiments with the well studied model organism *Acidobacterium capsulatus* at 1 % O_2 . While *A. capsulatus* also produced significant quantities of iso-diabolic acid (23 ± 2 %), we found no evidence of tetraether production at low O_2 (Table S-1). This suggests that brGDGT production at low O_2 is either not a universal trait among Acidobacteria, or that biosynthetic activation thresholds differ between organisms. Future genetic work with *E. aggregans* and additional culturing work with other Acidobacteria will help establish how widespread the genes for brGDGT production are in this phylum and under what conditions brGDGT biosynthesis occurs.

Palaeoclimate proxies. Although *E. aggregans* does not make a sufficiently large number of different tetraethers to test the mechanistic links between physiology and brGDGT-based

climate proxies such as MBT and CBT, the results presented here show clearly that O_2 limitation can be a trigger for brGDGT production. Because modern environmental calibrations rely on the relative distribution of brGDGTs in environmental samples, they are susceptible to abundance changes in any one structure. Although calibration data have demonstrated strong relationships between brGDGT distributions and environmental conditions like temperature and pH, dissolved O_2 measurements are limited in existing calibration data (Raberg *et al.*, 2021). It is therefore unknown whether this variable is implicitly captured or mostly unaccounted for in modern calibrations. Recent environmental observations indicate that the abundance of other brGDGTs not detected in this study (e.g., brGDGT IIa and IIIa) may also respond to redox conditions (Weber *et al.*, 2018; Yao *et al.*, 2020). Holocene palaeoclimate records from the Arctic show a decoupling between known temperature trends and brGDGT-inferred temperatures (e.g., Kusch *et al.*, 2019) suggesting an alternative overriding environmental control that is

unaccounted for. Our results suggest that for at least one source organism dissolved oxygen is the primary gradient to which brGDGT biosynthesis responds.

Conclusions

Our results indicate that O₂ availability controls biosynthesis of branched tetraethers by *E. aggregans*, with low O₂ required for production. This is the first confirmed organism to consistently produce significant quantities of multiple brGDGTs, thus opening the door to rigorous laboratory examination to elucidate the biosynthetic pathways and biological function of these enigmatic lipids. The identification of the enzymes involved in the synthesis of brGDGTs in *E. aggregans* will aid in the identification of other bacterial species that produce brGDGTs and help uncover the effect that O₂ limitation may have on brGDGT biosynthesis and palaeoclimate proxies.

Author Contributions

TAH, ADY and SHK designed the research. TAH, JMM, ADY, JD and ND performed the research. TAH and SHK analysed the data. TAH, JMM and SHK wrote the paper.

Acknowledgments

This research was supported by the Department of Geological Sciences at the University of Colorado Boulder and an NSF CAREER grant (EAR1945484) to SHK. We would like to thank J. Raberg and J. Sepúlveda for their analytical support and generous feedback.

Editor: Tanja Bosak

Additional Information

Supplementary Information accompanies this letter at <https://www.geochemicalperspectivesletters.org/article2132>.



© 2021 The Authors. This work is distributed under the Creative Commons Attribution 4.0 License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Additional information is available at <http://www.geochemicalperspectivesletters.org/copyright-and-permissions>.

Cite this letter as: Halamka, T.A., McFarlin, J.M., Younkin, A.D., Depoy, J., Dildar, N., Kopf, S.H. (2021) Oxygen limitation can trigger the production of branched GDGTs in culture. *Geochem. Persp. Let.* 19, 36–39.

References

- BOYER, G.M., SCHUBOTZ, F., SUMMONS, R.E., WOODS, J., SHOCK, E.L. (2020) Carbon Oxidation State in Microbial Polar Lipids Suggests Adaptation to Hot Spring Temperature and Redox Gradients. *Frontiers in Microbiology* 11, 229.
- DE JONGE, C., HOPMANS, E.C., ZELL, C.I., KIM, J.-H., SCHOUTEN, S., SINNINGHE DAMSTÉ, J.S. (2014) Occurrence and abundance of 6-methyl branched glycerol dialkyl glycerol tetraethers in soils: Implications for palaeoclimate reconstruction. *Geochimica et Cosmochimica Acta* 141, 97–112.
- EICHORST, S.A., TROJAN, D., ROUX, S., HERBOLD, C., RAITTE, T., WOEBKEN, D. (2018) Genomic insights into the Acidobacteriia reveal strategies for their success in terrestrial environments. *Environmental Microbiology* 20, 1041–1063.
- KOLTER, R., GREENBERG, E.P. (2006) The superficial life of microbes. *Nature* 441, 300–302.
- KUSCH, S., BENNIKE, O., WAGNER, B., LENZ, M., STEFFEN, I., RETHMEYER, J. (2019) Holocene environmental history in high-Arctic North Greenland revealed by a combined biomarker and macrofossil approach. *Boreas* 48, 273–286.
- LINCOLN, S.A., BRADLEY, A.S., NEWMAN, S.A., SUMMONS, R.E. (2013) Organic Geochemistry Archaeal and bacterial glycerol dialkyl glycerol tetraether lipids in chimneys of the Lost City Hydrothermal Field. *Organic Geochemistry* 60, 45–53.
- LIU, X.-L., ZHU, C., WAKEHAM, S.G., HINRICH, K.-U. (2014) In situ production of branched glycerol dialkyl glycerol tetraethers in anoxic marine water columns. *Marine Chemistry* 166, 1–8.
- LIU, X.-L., DE SANTIAGO TORIO, A., BOSAK, T., SUMMONS, R.E. (2016) Novel archaeal tetraether lipids with a cyclohexyl ring identified in Fayetteville Green Lake, NY, and other sulfidic lacustrine settings. *Rapid Communications in Mass Spectrometry* 30, 1197–1205.
- MARTÍNEZ-SOSA, P., TIERNEY, J.E. (2019) Lacustrine brGDGT response to microcosm and mesocosm incubations. *Organic Geochemistry* 127, 12–22.
- NAAFS, B.D.A., GALLEGO-SALA, A.V., INGLIS, G.N., PANCOST, R.D. (2017) Refining the global branched glycerol dialkyl glycerol tetraether (brGDGT) soil temperature calibration. *Organic Geochemistry* 106, 48–56.
- PETERSE, F., VAN DER MEER, J., SCHOUTEN, S., WEIJERS, J.W.H., FIERER, N., JACKSON, R.B., KIM, J.-H., SINNINGHE DAMSTÉ, J.S. (2012) Revised calibration of the MBT-CBT paleotemperature proxy based on branched tetraether membrane lipids in surface soils. *Geochimica et Cosmochimica Acta* 96, 215–229.
- PITCHER, R.S., WATMOUGH, N.J. (2004) The bacterial cytochrome cbb3 oxidases. *Biochimica et Biophysica Acta* 1655, 388–399.
- QIN, W., CARLSON, L.T., ARMBRUST, E.V., DEVOI, A.H., MOFFETT, J.W., STAHL, D.A., INGALLS, A.E. (2015) Confounding effects of oxygen and temperature on the TEX86 signature of marine Thaumarchaeota. *Proceedings of the National Academy of Sciences* 112, 10979–10984.
- RABERG, J.H., HARNING, D.J., CRUMP, S.E., DE WET, G., BLUMM, A., KOPF, S., GEIRSDÓTTIR, Á., MILLER, G.H., SEPÚLVEDA, J. (2021) Revised fractional abundances and warm-season temperatures substantially improve brGDGT calibrations in lake sediments. *Biogeosciences Discussions* 2021, 1–36.
- SINNINGHE DAMSTÉ, J.S., RIJSTRA, W.I.C., HOPMANS, E.C., WEIJERS, J.W.H., FOESEL, B.U., OVERMANN, J., DEDYSH, S.N. (2011) 13,16-Dimethyl Octacosanedioic Acid (iso-Diabolic Acid), a Common Membrane-Spanning Lipid of Acidobacteria Subdivisions 1 and 3. *Applied and Environmental Microbiology* 77, 4147–4154.
- SINNINGHE DAMSTÉ, J.S., RIJSTRA, W.I.C., FOESEL, B.U., HUBER, K.J., OVERMANN, J., NAKAGAWA, S., KIM, J.J., DUNFIELD, P.F., DEDYSH, S.N., VILLANUEVA, L. (2018) An overview of the occurrence of ether- and ester-linked isodiabolic acid membrane lipids in microbial cultures of the Acidobacteria: Implications for brGDGT paleoproxies for temperature and pH. *Organic Geochemistry* 124, 63–76.
- VAN DE VOSSENBERG, J.L.C.M., DRIESSEN, A.J.M., KONINGS, W.N. (1998) The essence of being extremophilic: The role of the unique archaeal membrane lipids. *Extremophiles* 2, 163–170.
- WANG, M., ZHENG, Z., ZONG, Y., MAN, M., TIAN, L. (2019) Distributions of soil branched glycerol dialkyl glycerol tetraethers from different climate regions of China. *Scientific Reports* 9, 1–8.
- WEBER, Y., DAMSTÉ, J.S.S., ZOPFI, J., DE JONGE, C., GILLI, A., SCHUBERT, C.J., LEFORI, F., LEHMANN, M.F., NIEMANN, H. (2018) Redox-dependent niche differentiation provides evidence for multiple bacterial sources of glycerol tetraether lipids in lakes. *Proceedings of the National Academy of Sciences* 115, 10926–10931.
- WEIJERS, J.W.H., SCHOUTEN, S., VAN DEN DONKER, J.C., HOPMANS, E.C., SINNINGHE DAMSTÉ, J.S. (2007) Environmental controls on bacterial tetraether membrane lipid distribution in soils. *Geochimica et Cosmochimica Acta* 71, 703–713.
- WESSEL, A.K., ARSHAD, T.A., FITZPATRICK, M., CONNELL, J.L., BONNECAZE, R.T., SHEAR, J.B., WHITELEY, M. (2014) Oxygen limitation within a bacterial aggregate. *mBio* 5, 1–9.
- YAO, Y., ZHAO, J., VACHULA, R.S., WERNE, J.P., WU, J., SONG, X., HUANG, Y. (2020) Correlation between the ratio of 5-methyl hexamethylated to pentamethylated branched GDGTs (HP5) and water depth reflects redox variations in stratified lakes. *Organic Geochemistry* 147, 104076.
- ZENG, Z., LIU, X.L., FARLEY, K.R., WEI, J.H., METCALF, W.W., SUMMONS, R.E., WELANDER, P.V. (2019) GDGT cyclization proteins identify the dominant archaeal sources of tetraether lipids in the ocean. *Proceedings of the National Academy of Sciences* 116, 22505–22511.

