

Residual forearc peridotites recording the early magmatic stage of subduction initiation


I. Nishio^{1,2*} , N. Akizawa¹, T. Ishii³, K. Itano⁴, A. Tamura⁵, T. Morishita^{5,6}



<https://doi.org/10.7185/geochemlet.2624>

Abstract

Residual forearc peridotites of early basaltic magmatism at subduction initiation



Subduction initiation is fundamental to plate tectonics. Early basaltic magmatism preceding boninitic magmatism is key to understanding the onset and mechanism of subduction initiation. However, the mantle processes during the early basaltic magmatism remain poorly understood due to the scarcity of corresponding residual peridotites. To find the residual peridotites from the early stage of subduction initiation, we compiled spinel data and examined peridotites from the Izu-Bonin-Mariana and Tonga forearc regions that contain some of the lowest spinel Cr/Al ratios. These samples present small amounts of pargasitic amphibole and light rare earth elements depleted patterns in their clinopyroxenes and amphiboles. These features differ from depleted forearc peridotites associated with boninitic magma, abyssal peridotites, and backarc peridotites, but resemble Iherzolites from the Oman ophiolite. Melting models can reproduce the compositions of clinopyroxene and the corresponding early basalts for each region. We suggest that the studied forearc peridotite samples are residues after early basaltic magma extractions, which potentially involved a common process of near fractional melting under moist conditions.

Received 12 January 2026 | Accepted 18 May 2026 | Published 30 June 2026

Introduction

Subduction initiation is a crucial process for the onset of plate tectonics and arises from a complex interplay between internal and external plate forces, plate structure, and physicochemical properties of the mantle (Stern and Gerya, 2018; Maunder *et al.*, 2020). Izu-Bonin-Mariana (IBM) and Tonga forearc regions in the western Pacific are the “modern” analogues (Fig. S-1), where tectonic configurations have been constrained by the studies on magmatic rocks and geophysical observation. Subduction initiation occurred around 50 Ma in both regions (Cramer *et al.*, 2020). In the IBM region, the Pacific Plate started to subduct beneath the Proto-Philippine Sea Plate along a pre-existing fracture zone (Ishizuka *et al.*, 2011). Following the onset of subduction initiation, the slab sinking induced a counter asthenospheric mantle flow, which drove seafloor spreading and generated an early mid-ocean ridge basalt (MORB)-like basalt, known as forearc basalt (FAB) (Reagan *et al.*, 2010). As the slab descended further, it released fluids that triggered melting of the residual mantle *via* slab-derived fluid influx, which in turn resulted in boninitic magmatism (Ishizuka *et al.*, 2020). In the Tonga region, subduction initiation occurred when the Pacific plate started to subduct beneath the Australian plate (Cramer *et al.*, 2020). This was triggered by the collision of the Papuan peninsula with the trench of the New Caledonia subduction

zone (Meffre *et al.*, 2012). The basalt in the Tonga region, interpreted as a product of subduction initiation, is known as early arc tholeiite (EAT) (Todd *et al.*, 2012). The early basaltic magmatism, which can provide the earliest record of subduction initiation, is the key to understanding subduction initiation mechanism.

Forearc peridotites are expected to record the magmatic evolution associated with the subduction initiation (*e.g.*, Birner *et al.*, 2017). Recovered forearc peridotites are mostly depleted, associated with boninitic magmatism, characterised by high spinel Cr# [$= \text{Cr}/(\text{Cr} + \text{Al})$] (>0.6 ; Fig. 1) (*e.g.*, Looke and Snow, 2024). Rarely, forearc peridotites with low spinel Cr# experienced reaction with early basaltic melts have been reported (Morishita *et al.*, 2011; Birner *et al.*, 2017). However, the residual forearc peridotites that retain the information of early basaltic magma extraction have not yet been identified. This is most likely because the signatures of the early basaltic magmatism were overprinted by subsequent melt-rock reaction, boninitic magmatism, and later geological processes. Such residual peridotites are the crucial piece for constraining the mantle processes corresponding to the early stage of subduction initiation. We systematically examined spinel data from forearc peridotites to evaluate their compositional variation and re-investigated forearc peridotite samples with among the lowest spinel Cr# from the IBM and Tonga regions.

1. Graduate School of Advanced Science and Engineering, Hiroshima University, Hiroshima 739-8526, Japan
 2. Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen 1350, Denmark
 3. Center for Integrated Research and Education of Natural Hazards, Shizuoka University, Shizuoka 422-8529, Japan
 4. Submarine Resources Research Program, Institute for Earth and Material Sciences, Japan Agency for Marine-Earth Science and Technology, Kanagawa 237-0061, Japan
 5. School of Geosciences and Civil Engineering, Kanazawa University, Kanazawa 920-1192, Japan
 6. Volcanoes and Earth's Interior Research Center, Research Institute for Marine Geodynamics, Japan Agency for Marine-Earth Science and Technology, Kanagawa 237-0061, Japan
- * Corresponding author (email: iknishio@hiroshima-u.ac.jp)




Figure 1 Spinel Cr# [= Cr/(Cr + Al)] vs. Mg# = [Mg/(Mg + Fe)] plot. The areas represent the 90 % density contours for spinels from abyssal and forearc peridotites, respectively. Abyssal spinel data are from Warren (2016). References for spinel data from the IBM and Tonga forearc peridotites are listed in the Supplementary Information.

Results

Morishita *et al.* (2011) and Birner *et al.* (2017) identified peridotites with low spinel Cr# that record reactions with early basalts (~0.5 for IBM and 0.3 for Tonga, respectively). Because melt-rock reactions generally increase spinel Cr# (Pearce *et al.*, 2000), residual peridotites that experienced only partial melting and basaltic melt extraction are expected to have lower spinel Cr#. Such samples (Cr# < 0.4) account only for ~4 % of forearc spinel data but do occur (Fig. 1). Based on the Cr# thresholds (~0.5 for IBM and 0.3 for Tonga), we investigated five samples with low spinel Cr# (four from IBM and one from Tonga) (Figs. S-2, S-3). The samples from the IBM region were collected from Conical seamount during ODP Leg 125 (Ishii *et al.*, 1992). IBM harzburgite sample (ODP125-778A-003R-CC-4-7) has spinel Cr# (0.38 ± 0.03 , average $\pm 2\sigma$) at the low end of the forearc peridotite

field (Fig. 1). The other three samples show spinel Cr# (average 0.45, 0.52, and 0.57), which are more similar to the Cr# threshold for IBM (~0.5) (Morishita *et al.*, 2011). For the Tonga region, we were able to investigate a lherzolite sample (BMRG08-111-3-6) with the lowest spinel Cr# (0.19 ± 0.03 ; Fig. 1) well below the threshold (0.3) for Tonga (Birner *et al.*, 2017). This sample was collected during the 1996 Boomerang Leg 8 cruise from the northern part of the Tonga forearc.

Rare earth elements (REEs) in clinopyroxenes from the lowest Cr# spinel-bearing IBM harzburgite and the Tonga lherzolite are characterised by light REE (LREE) depleted patterns with relatively high heavy REE (HREE) contents (Fig. 2). In the IBM harzburgite, LREE contents in clinopyroxenes are lower than detection limits, *i.e.* LREE depleted. The LREE depleted patterns of clinopyroxenes from these samples contrast with those from depleted forearc peridotites, which display high LREE/HREE ratios and low HREE contents (Day and Brown, 2021; Loocke and Snow, 2024). Clinopyroxenes from the IBM harzburgite show comparable REE contents to those from the LREE depleted residual abyssal peridotites formed after extensive partial melting and melt extraction. Clinopyroxenes from the Tonga lherzolite show similar REE patterns to those from general abyssal peridotites (Warren, 2016). Clinopyroxenes from the IBM samples with slightly higher spinel Cr# (>0.45) show lower HREE contents with detectable contents of LREE, which share the characteristics of depleted forearc peridotites.

Pargasitic amphiboles (<0.1 vol. %), are observed in the IBM harzburgite and the Tonga lherzolite with the lowest Cr#, where they are surrounded by orthopyroxene and/or adjacent to clinopyroxene (Fig. 3a,b). Amphiboles are often found in forearc and backarc peridotites (Morishita *et al.*, 2011; Akizawa *et al.*, 2021) but are rare in abyssal peridotites (Cipriani *et al.*, 2009). Amphiboles also occur in basal lherzolites with low spinel Cr# (~0.2) of the supra-subduction zone Oman ophiolite (Prigent *et al.*, 2018). Pargasitic amphiboles show relatively high TiO₂ (IBM, 0.45 wt. %; Tonga, 0.75 wt. %) and low K₂O contents (<~0.1 wt. %) (Fig. 3c), which resemble those from abyssal peridotites and from Oman basal lherzolites. Amphiboles from




Figure 2 Chondrite normalised REE patterns of clinopyroxene from (a) the IBM forearc peridotites and (b) the Tonga forearc peridotites. Cr# indicates the value of the coexisting spinel. References for clinopyroxene data of IBM peridotites, Tonga peridotites, and fertile and depleted abyssal peridotites are listed in the SI.




Figure 3 Back scatter electron images of (a) the IBM harzburgite (ODP125-778A-003R-CC-4-7) and (b) the Tonga lherzolite (BMRG08-111-3-6). Ol, olivine; Opx, orthopyroxene; Cpx, clinopyroxene; and Prg, pargasite. (c) TiO₂-K₂O discrimination plot and (d) chondrite normalised REE patterns for amphibole from the IBM harzburgite and the Tonga lherzolite, compared with data from forearc, ultra-depleted, backarc, and abyssal peridotites, as well as Oman basal lherzolites. Areas for mid-ocean ridge basalt (MORB), backarc basin basalt (BABB), forearc basalt and early arc tholeiite (FAB-EAT), and boninite are shown in (c). References for amphibole data are provided in the SI.

backarc and depleted forearc peridotites show distinct compositions: backarc amphiboles display variable and higher TiO₂ and K₂O contents, whereas amphiboles from depleted peridotites are characterised by low TiO₂ and high K₂O contents.

Pargasitic amphiboles in the Tonga lherzolite show LREE depleted patterns similar to those of the coexisting clinopyroxenes, and to amphiboles from Oman basal lherzolites (Fig. 3d). We could not analyse trace element compositions of pargasite in the IBM harzburgite due to its small grain size. Notably, amphiboles and clinopyroxenes from one IBM sample exhibit LREE contents below detection limits (Figs. 2a, 3d; Ichiyama *et al.*, 2021). Such LREE depleted patterns differ from those of backarc peridotites (Gong *et al.*, 2022), abyssal peridotites (Cipriani *et al.*, 2009), depleted peridotites in relation to boninitic magmas (Nishio *et al.*, 2023), and IBM dunites formed by reaction with FAB (Morishita *et al.*, 2011).

Sample information, analytical methods, and mineral major and trace element compositions are provided in the SI and SI Tables.

Discussion

IBM harzburgite ODP125-778A-003R-CC-4-7 and Tonga lherzolite BMRG08-111-3-6 are characterised by clinopyroxenes with LREE depleted patterns and relatively high HREE contents, and low spinel Cr# (Figs. 1, 2). These characteristics align them with residual abyssal peridotites that experienced partial melting and basaltic melt extraction without significant melt-rock reaction (Warren, 2016). Clinopyroxenes from the IBM samples with slightly high spinel Cr# (0.45, 0.52, and 0.57) show lower HREE and detectable LREE contents. Reactions with melts or fluids generally increase LREE contents in residues (Birner *et al.*, 2017). In our compiled forearc spinel data set ($n = 287$), 4 % of forearc peridotites exhibit low spinel Cr# (<0.4) and one IBM peridotite exhibits LREE depleted clinopyroxene patterns (Ichiyama *et al.*, 2021). The rare forearc peridotite samples with low spinel Cr# and LREE depleted clinopyroxene patterns represent potential mantle residues that preserve a record of partial melting during early basaltic magmatism of subduction initiation, without significant overprinting by melt-rock reactions.

Although the number of available samples is very limited and sampling bias cannot be ruled out, these samples nevertheless provide crucial direct evidence for early basaltic magmatism.

We therefore examined the samples (IBM, ODP125-778A-003R-CC-4-7; Tonga, BMRG08-111-3-6) to investigate melting processes and their link to early basaltic magmatism using an open system melting model (Ozawa, 2001). We conducted influx-free melting models using a depleted MORB mantle (DMM) source or a depleted DMM (DDMM) source (Workman and Hart, 2005). We varied two key parameters: the critical melt fraction (a_c : 0–2 %) at which melt starts to segregate, and the degree of melting (F : 0–25 %), with melting initiated in the garnet peridotite stability field (F_{grt} : 0–10 %) prior to the spinel peridotite stability fields. We compared the modelled clinopyroxene compositions ($n = \sim 20,000$) with those of the studied samples corrected for subsolidus exchange, respectively (Figs. 4 and S-4–S-7). The modelled accumulated melt compositions were compared with FAB from the IBM region (Ishizuka *et al.*, 2011; Shervais *et al.*, 2019) and EAT from the Tonga region (Todd *et al.*, 2012), respectively.

Details of model set up, subsolidus correction, best fit model estimation, and model results are provided in the SI.

Our models show that REE contents in clinopyroxene decrease with increasing F . At a given F , increasing a_c suppresses LREE depletion, whereas increasing F_{grt} suppresses HREE depletion in clinopyroxene of spinel peridotite (Fig. S-5). At given F (>10 %), accumulated melt shows similar REE patterns regardless of a_c and F_{grt} (Fig. S-6). The modal Ce contents in accumulated melts are 2.98 ppm for models derived from the DDMM source and 3.86 ppm for those derived from the DMM source (Fig. S-7), reflecting slightly lower LREE contents in melts from the DDMM source.

Best fit models based on reduced chi square successfully reproduce the REE patterns of both clinopyroxene and basalt in each region (Figs. 4, S-8). For the IBM samples, the best fits correspond to $F = 16.8 \pm 2.1$ % (2σ) with $a_c = 0.4 \pm 0.6$ % and $F_{\text{grt}} = 2.9 \pm 3.8$ %. For the Tonga samples, the best fits correspond to $F = 7.4 \pm 0.1$ % with $a_c = 1$ % and $F_{\text{grt}} = 4$ %. Among IBM best fit models, 50.9 % adopt the DDMM source and 40.9 % the DMM source, whereas all Tonga best fit models are derived from the DMM source. Very low a_c (≤ 1 %) indicates that the melting conditions recorded in the studied forearc peridotites are near fractional melting, which is similar to the melting conditions beneath mid-ocean ridges (*e.g.*, Warren, 2016).

Given the similarity between the studied forearc peridotites and abyssal peridotites, there is a possibility that the studied samples might be remnants of abyssal peridotites that did not experience melting during subduction initiation. However, the studied samples contain minor amounts of pargasitic amphibole (Fig. 3) that are a common mineral phase in the forearc peridotites but not in the abyssal peridotites (Cipriani *et al.*, 2009; Morishita *et al.*, 2011). High TiO_2 and low K_2O contents in amphiboles from the studied samples support an origin related to early basalts (FAB and EAT) or MORB (Fig. 3c). LREE depleted patterns in amphiboles and clinopyroxenes from our studied samples and the IBM sample reported by Ichiyama *et al.* (2021) further contrast with those of amphiboles from abyssal peridotites, backarc peridotites, and ultra-depleted peridotites in relation to boninitic magmatism (Fig. 3d). Such LREE depleted clinopyroxenes and amphiboles have been reported from Oman basal lherzolites, where depleted harzburgites associated with boninites also occur (Prigent *et al.*, 2018). We therefore interpret the studied forearc peridotites as residues formed by partial melting and melt extraction during early basaltic magmatism at subduction initiation.




Figure 4 Chondrite normalised REE patterns: (a,b) IBM and (c–e) Tonga, showing (a,c) modelled accumulated melt compared with FAB and EAT, (b,d) modelled clinopyroxene compared with subsolidus-corrected clinopyroxene in IBM harzburgite ODP125-778A-003R-CC-4-7 and Tonga lherzolite BMRG08-111-3-6, and (e) modelled amphibole compared with amphiboles in the Tonga lherzolite. Sample symbols represent average values with ± 2 standard deviations. All models are calculated at fixed melting degree (F), critical melt fraction (a_c), and source based on the best fit models (Fig. S-8), indicated at the top of each column, with colour denoting influx rate (β).

We further performed slab derived fluid (Li *et al.*, 2013; Shervais *et al.*, 2019) influx open system melting models to evaluate the effect of fluid infiltration and amphibole formation (Figs. 4 and S-9–S-11). Model settings are based on the best fit conditions constrained above, with the influx rate (β) varied from 0 to 1 and include amphibole crystallisation from a small fraction of trapped melt (0.1 %). At the best fit F (IBM, 16.8 %; Tonga, 7.4 %), increasing β suppresses LREE depletion in clinopyroxene (Fig. 4b,d), and amphibole shows similar REE behaviour (Fig. 4e). The effect of fluid infiltration on the accumulated melt is comparatively minor (Fig. 4a,c).

To reproduce the REE patterns of the studied clinopyroxene, β was required to be <0.0001 for the IBM samples to match LREE contents below detection limits, and <0.01 for the Tonga samples (Fig. 4b,d). These models simultaneously reproduced REE patterns of FAB and EAT, respectively (Fig. 4a,c), and capture the LREE depleted patterns of the Tonga amphibole (Fig. 4e). The very low a_c (≤ 1 %) and β (<0.01), with the small trapped melt fraction (~ 0.1 %) also agree with the minor amounts of amphibole in the samples. Amphiboles in the studied samples may have formed during the early basaltic magmatic stage, through a combination of reactions with infiltrating fluids (Prigent *et al.*, 2018) at very low β , and crystallisation from a small fraction of trapped melts. Although the source of the fluid remains uncertain, our results imply that the melting conditions were not dry. “Moist” conditions have been proposed for the FAB formation based on their depleted REE patterns with slightly higher H_2O/Ce ratios compared to MORB (Brounce *et al.*, 2021), and lower TiO_2 contents at similar MgO contents relative to MORB (MacLeod *et al.*, 2013). Our findings from mantle peridotite samples are consistent with those of FAB studies.

Our best fit models showed that the Tonga samples can be reproduced by models with the DMM, whereas the IBM samples can be reproduced by models using either the DMM or the DDMM (Figs. 4, S-8). Accumulated melts in the IBM best fit models are slightly more LREE enriched when derived from the DMM compared to the DDMM, as reflected in the modal Ce contents (Fig. S-7). Studies on FAB have also suggested a source depleted by ~ 0.5 – 2 % relative to the DMM (Umino *et al.*, 2018; Yagodinski *et al.*, 2018; Shervais *et al.*, 2019). This degree of prior depletion is consistent with the difference between the DMM and DDMM sources (Workman and Hart, 2005). Therefore, a source more depleted than the DMM, such as the DDMM may be more consistent with the IBM samples. The compiled forearc spinel data set shows that Cr# above 0.8 occur in the IBM, but not in the Tonga (Fig. 1), in agreement with the higher best fit F in the IBM (16.8 %) compared to the Tonga (7.4 %). Although further study is necessary, our model results and spinel Cr# systematics indicate potential regional variability, in agreement with studies of early basalts (Arculus *et al.*, 2019), which potentially reflect differences such as the subduction initiation style and source mantle composition.

Conclusions

We examined the least depleted harzburgites from the IBM forearc and a lherzolite from the Tonga forearc among the rare ~ 4 % population among forearc peridotites. They show low spinel Cr#, and LREE depleted patterns in clinopyroxenes and amphiboles. These characteristics differ from the typical depleted forearc peridotites after boninitic magma extraction. Our open system melting models reproduced the REE contents of the studied clinopyroxenes and early basalts, and amphiboles of each region. We identified the IBM and Tonga peridotites as residual peridotites after the extraction of early basaltic magma during subduction initiation. The LREE depleted patterns of clinopyroxenes

and amphiboles suggest that the IBM and Tonga peridotites may have formed through near fractional melting under moist conditions. These forearc peridotites are very rare, but essential to understand the controlling factors of subduction initiation such as source mantle compositions and style of subduction initiation style.

Data Availability

All data and Supplementary Information are available in online.

Acknowledgments

We thank P. Waterton and K. Ozawa for their help with the modelling and best fit estimation, and T. Mizukami, and Y. Ichiyama for discussions, and H. Yamashita for providing the samples. We thank J. Pearce, K.Y. Lin, and J. Hermann for reviews of a previous version of this manuscript. We thank V. Basch, an anonymous reviewer, and Editor R.O.C. Fonseca for their constructive comments, which improved the paper.

Editor: Raul O.C. Fonseca

Funding

This study was supported by Japan Society for the Promotion of Science (no. 24JK0076 and no. 23K25963).

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Additional Information

Supplementary Information accompanies this letter at <https://www.geochemicalperspectivesletters.org/article2624>.



© 2026 The Authors. This work is distributed under the Creative Commons Attribution Non-Commercial No-Derivatives 4.0

License, which permits unrestricted distribution provided the original author and source are credited. The material may not be adapted (remixed, transformed or built upon) or used for commercial purposes without written permission from the author. Additional information is available at <https://www.geochemicalperspectivesletters.org/copyright-and-permissions>.

Cite this letter as: Nishio, I., Akizawa, N., Ishii, T., Itano, K., Tamura, A., Morishita, T. (2026) Residual forearc peridotites recording the early magmatic stage of subduction initiation. *Geochim. Persp. Let.* 41, 1–6. <https://doi.org/10.7185/geochemlet.2624>

References

- AKIZAWA, N., OHARA, Y., OKINO, K., ISHIZUKA, O., YAMASHITA, H., MACHIDA, S., SANFILIPPO, A., BASCH, V., SNOW, J.E., SEN, A., HIRAUCHI, K.-I., MICHIBAYASHI, K., HARIGANE, Y., FUJII, M., ASANUMA, H., HIRATA, T. (2021) Geochemical characteristics of back-arc basin lower crust and upper mantle at final spreading stage of Shikoku Basin: an example of Mado Megamullion. *Progress in Earth and Planetary Science* 8, 65. <https://doi.org/10.1186/s40645-021-00454-3>

- ARCULUS, R.J., GURNIS, M., ISHIZUKA, O., REAGAN, M.K., PEARCE, J.A., SUTHERLAND, R. (2019) How to create new subduction zones: A global perspective. *Oceanography* 32, 160–174. <https://doi.org/10.5670/oceanog.2019.140>
- BIRNER, S.K., WARREN, J.M., COTTELL, E., DAVIS, F.A., KELLEY, K.A., FALLOON, T.J. (2017) Forearc Peridotites from Tonga Record Heterogeneous Oxidation of the Mantle following Subduction Initiation. *Journal of Petrology* 58, 1755–1780. <https://doi.org/10.1093/ptrology/egx072>
- BROUNCE, M., REAGAN, M.K., KELLEY, K.A., COTTELL, E., SHIMIZU, K., ALMEEV, R. (2021) Covariation of Slab Tracers, Volatiles, and Oxidation During Subduction Initiation. *Geochemistry, Geophysics, Geosystems* 22, e2021GC009823. <https://doi.org/10.1029/2021GC009823>
- CIPRIANI, A., BONATTI, E., SEYLER, M., BRUECKNER, H.K., BRUNELLI, D., DALLAI, L., HEMMING, S.R., LIGI, M., OTTOLINI, L., TURRIN, B.D. (2009) A 19 to 17 Ma amagmatic extension event at the Mid-Atlantic Ridge: Ultramafic mylonites from the Vema Lithospheric Section. *Geochemistry, Geophysics, Geosystems* 10, Q10011. <https://doi.org/10.1029/2009GC002534>
- CRAMER, F., MAGNI, V., DOMEIER, M., SHEPHARD, G.E., CHOTALIA, K., COOPER, G., EAKIN, C.M., GRIMA, A.G., GÜNER, D., KIRÁLY, Á., MULYUKOVA, E., PETERS, K., ROBERT, B., THIELMANN, M. (2020) A transdisciplinary and community-driven database to unravel subduction zone initiation. *Nature Communications* 11, 3750. <https://doi.org/10.1038/s41467-020-17522-9>
- DAY, J.M.D., BROWN, D.B. (2021) Ancient Melt-Depletion in Fresh to Strongly Serpentinized Tonga Trench Peridotites. *Journal of Petrology* 62, egab088. <https://doi.org/10.1093/ptrology/egab088>
- GONG, X., TIAN, L., DONG, Y. (2022) Contrasting melt percolation and melt-rock reactions in the Parece Vela back-arc oceanic lithosphere, Philippine Sea: A mineralogical perspective. *Lithos* 422–423, 106727. <https://doi.org/10.1016/j.lithos.2022.106727>
- ICHIYAMA, Y., TSUJIMORI, T., FRYER, P., MICHIBAYASHI, K., TAMURA, A., MORISHITA, T. (2021) Temporal and spatial mineralogical changes in clasts from Mariana serpentinite mud volcanoes: Cooling of the hot forearc-mantle at subduction initiation. *Lithos* 384–385, 105941. <https://doi.org/10.1016/j.lithos.2020.105941>
- ISHII, T., ROBINSON, P.T., MAEKAWA, H., FISKE, R. (1992) Petrological studies of peridotites from diapiric serpentinite seamounts in the Izu-Ogasawara-Mariana Forearc, Leg 125. In: FRYER, P., PEARCE, J.A., STOKING, L.B., ALI, J.R., ARCULUS, R., et al. (Eds.) *Proceedings of the Ocean Drilling Program, Scientific Results*, 125, Ocean Drilling Program, College Station, TX, 445–485. <https://doi.org/10.2973/odp.proc.sr.125.129.1992>
- ISHIZUKA, O., TANI, K., REAGAN, M.K., KANAYAMA, K., UMINO, S., HARIGANE, Y., SAKAMOTO, I., MIYAJIMA, Y., YUASA, M., DUNKLEY, D.J. (2011) The timescales of subduction initiation and subsequent evolution of an oceanic island arc. *Earth and Planetary Science Letters* 306, 229–240. <https://doi.org/10.1016/j.epsl.2011.04.006>
- ISHIZUKA, O., TAYLOR, R.N., UMINO, S., KANAYAMA, K. (2020) Geochemical Evolution of Arc and Slab Following Subduction Initiation: a Record from the Bonin Islands, Japan. *Journal of Petrology* 61, egaa050. <https://doi.org/10.1093/ptrology/egaa050>
- LI, Y.-B., KIMURA, J.-I., MACHIDA, S., ISHII, T., ISHIWATARI, A., MARUYAMA, S., QIU, H.-N., ISHIKAWA, T., KATO, Y., HARAGUCHI, S., TAKAHATA, N., HIRAHARA, Y., MIYAZAKI, T. (2013) High-Mg Adakite and Low-Ca Boninite from a Bonin Fore-arc Seamount: Implications for the Reaction between Slab Melts and Depleted Mantle. *Journal of Petrology* 54, 1149–1175. <https://doi.org/10.1093/ptrology/egt008>
- LOOCKE, M.P., SNOW, J.E. (2024) Arc Foundations and Subduction Initiation: Insights into the Magmatic Evolution of the Lower Crust/Upper Mantle of the Izu-Bonin Forearc during Subduction Initiation. *Journal of Petrology* 65, egae038. <https://doi.org/10.1093/ptrology/egae038>
- MACLEOD, C.J., LISSNERBERG, C.J., BIBBY, L.E. (2013) “Moist MORB” axial magmatism in the Oman ophiolite: The evidence against a mid-ocean ridge origin. *Geology* 41, 459–462. <https://doi.org/10.1130/G33904.1>
- MAUNDER, B., PRYTULAK, J., GOES, S., REAGAN, M. (2020) Rapid subduction initiation and magmatism in the Western Pacific driven by internal vertical forces. *Nature Communications* 11, 1874. <https://doi.org/10.1038/s41467-020-15737-4>
- MEFFRE, S., FALLOON, T.J., CRAWFORD, T.J., HOERNLE, K., HAUFF, F., DUNCAN, R.A., BLOOMER, S.H., WRIGHT, D.J. (2012) Basalts erupted along the Tongan fore arc during subduction initiation: Evidence from geochronology of dredged rocks from the Tonga fore arc and trench. *Geochemistry, Geophysics, Geosystems* 13, Q12003. <https://doi.org/10.1029/2012GC004335>
- MORISHITA, T., TANI, K., SHUKUNO, H., HARIGANE, Y., TAMURA, A., KUMAGAI, H., HELLEBRAND, E. (2011) Diversity of melt conduits in the Izu-Bonin-Mariana forearc mantle: Implications for the earliest stage of arc magmatism. *Geology* 39, 411–414. <https://doi.org/10.1130/G31706.1>
- NISHIO, I., MORISHITA, T., TAMURA, A., ITANO, K., TAKAMIZAWA, S., ICHYAMA, Y., ARAI, S., BARRETT, N., SZILAS, K. (2023) Formation of Ultra-Depleted Mantle Peridotites and Their Relationship With Boninitic Melts: An Example From the Kamuikotan Unit, Hokkaido, Japan. *Journal of Geophysical Research: Solid Earth* 128, e2022JB025066. <https://doi.org/10.1029/2022JB025066>
- OZAWA, K. (2001) Mass balance equations for open magmatic systems: Trace element behavior and its application to open system melting in the upper mantle. *Journal of Geophysical Research: Solid Earth* 106, 13407–13434. <https://doi.org/10.1029/2001JB900001>
- PEARCE, J.A., BARKER, P.F., EDWARDS, S.J., PARKINSON, I.J., LEAT, P.T. (2000) Geochemistry and tectonic significance of peridotites from the South Sandwich arc-basin system, South Atlantic. *Contributions to Mineralogy and Petrology* 139, 36–53. <https://doi.org/10.1007/s004100050572>
- PRIGENT, C., AGARD, P., GUILLLOT, S., GODARD, M., DUBACQ, B. (2018) Mantle Wedge (De)formation During Subduction Infancy: Evidence from the Base of the Semail Ophiolitic Mantle. *Journal of Petrology* 59, 2061–2092. <https://doi.org/10.1093/ptrology/egy090>
- REAGAN, M.K., ISHIZUKA, O., STERN, R.J., KELLEY, K.A., OHARA, Y., BLICHERT-TOFT, J., BLOOMER, S.H., CASH, J., FRYER, P., HANAN, B.B., HICKEY-VARGAS, R., ISHII, T., KIMURA, J.-I., PEATE, D.W., ROWE, M.C., WOODS, M. (2010) Fore-arc basalts and subduction initiation in the Izu-Bonin-Mariana system. *Geochemistry, Geophysics, Geosystems* 11, Q03X12. <https://doi.org/10.1029/2009GC002871>
- SHERVAIS, J.W., REAGAN, M., HAUGEN, E., ALMEEV, R.R., PEARCE, J.A., PRYTULAK, J., RYAN, J.G., WHATTAM, S.A., GODARD, M., CHAPMAN, T., LI, H., KURZ, W., NELSON, W.R., HEATON, D., KIRCHENBAUR, M., SHIMIZU, K., SAKUYAMA, T., LI, Y., VETTER, S.K. (2019) Magmatic Response to Subduction Initiation: Part 1. Fore-arc Basalts of the Izu-Bonin Arc From IODP Expedition 352. *Geochemistry, Geophysics, Geosystems* 20, 314–338. <https://doi.org/10.1029/2018GC007731>
- STERN, R.J., GERYA, T. (2018) Subduction initiation in nature and models: A review. *Tectonophysics* 746, 173–198. <https://doi.org/10.1016/j.tecto.2017.10.014>
- TODD, E., GILL, J.B., PEARCE, J.A. (2012) A variably enriched mantle wedge and contrasting melt types during arc stages following subduction initiation in Fiji and Tonga, southwest Pacific. *Earth and Planetary Science Letters* 335–336, 180–194. <https://doi.org/10.1016/j.epsl.2012.05.006>
- UMINO, S., KANAYAMA, K., KITAMURA, K., TAMURA, A., ISHIZUKA, O., SENDA, R., ARAI, S. (2018) Did boninite originate from the heterogeneous mantle with recycled ancient slab? *Island Arc* 27, e12221. <https://doi.org/10.1111/iar.12221>
- WARREN, J.M. (2016) Global variations in abyssal peridotite compositions. *Lithos* 248–251, 193–219. <https://doi.org/10.1016/j.lithos.2015.12.023>
- WORKMAN, R.K., HART, S.R. (2005) Major and trace element composition of the depleted MORB mantle (DMM). *Earth and Planetary Science Letters* 231, 53–72. <https://doi.org/10.1016/j.epsl.2004.12.005>
- YOGODZINSKI, G.M., BIZMIS, M., HICKEY-VARGAS, R., MCCARTHY, A., HOCKING, B.D., SAVOV, I.P., ISHIZUKA, O., ARCULUS, R. (2018) Implications of Eocene-age Philippine Sea and forearc basalts for initiation and early history of the Izu-Bonin-Mariana arc. *Geochimica et Cosmochimica Acta* 228, 136–156. <https://doi.org/10.1016/j.gca.2018.02.047>



Residual forearc peridotites recording the early magmatic stage of subduction initiation

I. Nishio, N. Akizawa, T. Ishii, K. Itano, A. Tamura, T. Morishita

Supplementary Information

The Supplementary Information includes:

- Samples and Analytical Methods
- Reconstructing Primary REE Contents in Clinopyroxene
- Influx-Free Open System Melting Models
- Best Fit Model Estimation
- Slab Derived Fluid Influx Open System Melting Models
- Compiled Forearc Spinel Data and Reference Data for Plots
- Tables S-1 to S-15
- Figures S-1 to S-12
- Supplementary Information References

Samples and Analytical Methods

The samples are four harzburgites (ODP125-778A-003R-CC-4-7, ODP125-779A-26R-02-47-50, ODP125-780C-10R-01-11-13, ODP125-780D-005X-01-0-4) from Conical seamount in the IBM forearc (Fig. S-1), which were collected during ODP Leg 125 Site 778A, 779A, 780C, and 780D (Fig. S-2) (Ishii *et al.*, 1992; Parkinson *et al.*, 1992; Parkinson and Pearce, 1998) and a lherzolite sample (BMRG08-111-3-6) from the northern part of the Tonga forearc, which was collected during the 1996 Boomerang Leg 8 cruise (Bloomer *et al.*, 1996; Wright *et al.*, 2000; Birner *et al.*, 2016, 2017). We use samples previously studied by Ishii *et al.* (1992) (IBM) and Birner *et al.* (2017) (Tonga). IBM harzburgite ODP125-778A-003R-CC-4-7 and Tonga lherzolite BMRG08-111-3-6 contain minor amounts of clinopyroxene (IBM: ~2 vol.%; Tonga: 9 vol.%). Plagioclase and garnet are absent in the samples, and the IBM samples are serpentinised.




Figure S-1 A map of the western Pacific. Arrows in the IBM and Tonga represent the sample locations.




Figure S-2 Photomicrographs of IBM harzburgite and Tonga lherzolite samples. **(a and b)** The harzburgite sample (ODP125-778A-003R-CC-4-7) from the Conical Seamount IBM forearc region. **(c and d)** The lherzolite sample (BMRG08-111-3-6) from the Northern Tonga forearc region. The red colour bar indicates 2 millimetres. (b) and (d) were taken under cross-polarized light.




Figure S-3 Mineral modes of the IBM harzburgite (ODP125-778A-003R-CC-4–7) and Tonga lherzolite (BMRG08-111-3-6). Comparison of mineral modes among the studied samples, residual abyssal peridotites (Warren, 2016), and depleted forearc peridotites (Parkinson and Pearce, 1998; Birner *et al.*, 2017; Loocke and Snow, 2024).

Major element compositions of minerals in peridotite samples were obtained using the JEOL JXA-8900R electron microprobe at the Atmosphere and Ocean Research Institute, University of Tokyo. Analytical conditions were set at an accelerating voltage of 15 kV, 12 nA probe current, and a 3 μm beam spot for the analyses. Natural and synthetic mineral standards were used for calibration, and ZAF correction was used for data reduction. In-house mineral standards (olivine, chromian spinel, diopside, and enstatite) were measured to monitor data quality (Akizawa *et al.*, 2021). We determined Fe^{3+} content in spinel using the method that initially treats Fe as Fe^{2+} and assumes any cation excess as Fe^{3+} beyond the three cations per four oxygens. Elemental distribution mapping of the thin section was performed using an accelerating voltage of 15 kV, a probe current of 80 nA, a focused probe with a 20 μm diameter, a step size of 30 μm , and a dwell time of 100 ms. Modal mineral abundances were determined using a supervised machine learning method implemented in XMapTools (Lanari *et al.*, 2014; Lanari and Tedeschi, 2025). To obtain representative mineral modes, we performed six repeated classifications, although potential bias introduced during thin section preparation could not be eliminated (Fig. S-3).

Trace element compositions of silicates were determined by a laser ablation-inductively coupled plasma-mass spectrometry (LA-ICP-MS) (New Wave Research UP-213 laser system and Agilent 7850 ICP-MS) at Kanazawa University. A laser beam diameter of 100 μm was used for orthopyroxene, amphibole, and olivine, and either 100 μm or 55 μm was used for clinopyroxene, depending on the size, with a repetition rate of 6 or 10 Hz and an energy density of 8 J/cm². The NIST 612 standard was used as the primary calibration standard and was analysed at the beginning of each batch ($n < 8$) with a linear drift correction applied within each analytical session. Standard glass NIST 612 was used for calibration with ²⁹Si as an internal standard, based on the Si content obtained by electron microprobe analysis (Longerich *et al.*, 1996; Pearce *et al.*, 1997; Jochum *et al.*, 2011). Details of analytical procedures and quality checks were provided in Sagawa *et al.* (2022) and Tamura *et al.* (2022).

Reconstructing Primary REE Contents in Clinopyroxene

The mineral compositions of peridotites commonly reflect subsolidus re-equilibration rather than compositions at solidus conditions (Witt-Eickschen and O'Neill, 2005; Liang *et al.*, 2013; Liang *et al.*, 2021; Lin *et al.*, 2023). We corrected the REE concentrations in clinopyroxene at a given temperature using modal abundances and partition coefficients (Eq. S-1). We did not consider contributions from olivine because of their very low concentrations and partition coefficients:

$$C_{cpx}^{corrected} = \frac{\omega_{cpx}^{measured} + \omega_{opx}^{measured} \times D_{cpx/opx}^{measured}}{\omega_{cpx}^{corrected} + \omega_{opx}^{corrected} \times D_{cpx/opx}^{corrected}} \times C_{cpx}^{measured} \quad (\text{Eq. S-1})$$

$C_{cpx}^{corrected}$ is the concentration of element i (REE) in clinopyroxene corrected for subsolidus re-equilibration at a given temperature. $\omega_{cpx}^{measured}$ and $\omega_{opx}^{measured}$ are analysed modal abundances of clinopyroxene and orthopyroxene. $D_{cpx/opx}^{measured}$ is the exchange coefficient between clinopyroxene and orthopyroxene at the sample temperature. $C_{cpx}^{measured}$ is the analysed concentration of element i in clinopyroxene. $\omega_{cpx}^{corrected}$ and $\omega_{opx}^{corrected}$ are corrected modal abundances of clinopyroxene and orthopyroxene. $D_{cpx/opx}^{corrected}$ is the exchange coefficient between clinopyroxene and orthopyroxene at a given temperature.

We estimated the exchange partition coefficient at a given temperature using the semi-empirical model (Witt-Eickschen and O'Neill, 2005) and the lattice-strain model (Sun and Liang, 2012; Yao *et al.*, 2012; Liang *et al.*, 2013; Sun and Liang, 2013). In the semi-empirical model of Witt-Eickschen and O'Neill (2005), the exchange partition coefficient for REE is described by a nonlinear regression equation:

$$\ln D_{cpx/opx} = a_0 + \frac{a_1 + a_2 \cdot N_{Na}^{cpx} + a_3 \cdot Cr_{spl}}{T_{opx}(1.5GPa)} \quad (\text{Eq. S-2}),$$

where: a_0 to a_3 are the regression coefficients. N_{Na}^{cpx} is the number of atoms of Na in clinopyroxene per formula unit of six oxygen. Cr_{spl} is the molar Cr/(Cr + Al) in spinel. $T_{opx}(1.5GPa)$ is the temperature estimated by Ca-in-orthopyroxene geothermometer at 1.5 GPa (Brey and Köhler, 1990). We used average values for each region: IBM harzburgite ($N_{Na}^{cpx} = 0.0156$, $Cr_{spl} = 0.378$) and Tonga lherzolite ($N_{Na}^{cpx} = 0.0345$, $Cr_{spl} = 0.195$). The Ca in orthopyroxene geothermometer often records lower temperatures than REE in two pyroxene for exhumed peridotite bodies that have experienced slow cooling, because Ca has a lower closure temperature than REE (Dygert and Liang, 2015). Applying the Ca in orthopyroxene geothermometer to our forearc peridotite samples would likely overestimate the extent of REE subsolidus exchange. Therefore, we used temperatures calculated from the REE-in two pyroxene geothermometer at 1.5 GPa (Liang *et al.*, 2013). The estimated temperatures were 919 ± 3 °C (1 σ) for the IBM harzburgite (ODP125-778A-003R-CC-4–7) and $931^\circ\text{C} \pm 31$ for the Tonga lherzolite (BMRG08-111-3-6).

For the lattice-strain model, we obtained partition coefficients using an Excel macro (Sun and Liang, 2012; Yao *et al.*, 2012; Liang *et al.*, 2013; Sun and Liang, 2013) following Lin *et al.* (2023). We used the average major compositions of clinopyroxene and orthopyroxene for each sample and used TiO₂ contents of primitive basalts (0.63 wt.% for FAB and 1.27 wt.% for EAT) with MgO > 7.5 wt.% (Ishizuka *et al.*, 2011; Todd *et al.*, 2012; Shervais *et al.*, 2019) and an H₂O content of 0.85 wt.% in FAB (Brounce *et al.*, 2021) to calculate the partition coefficient.

Using the exchange coefficients obtained by two different models, we corrected REE concentrations in clinopyroxene to temperature of 1000–1100 °C. At subsolidus temperatures, clinopyroxene can exsolve from orthopyroxene (Poldervaart and Hess, 1951; Arai, 1984). We recalculated the mineral modes at 1000–1100 °C, assuming that 5% of the orthopyroxene exsolved to clinopyroxene during the subsolidus conditions (Akizawa *et al.*, 2016). For a semi-empirical model, Pr, Tb, and Tm contents were interpolated from adjacent elements on the chondrite-normalized REE pattern. To account for uncertainty, we repeated the calculations by combining multiple parameters. We considered two sets of exchange coefficients (D) and three temperature estimates from the REE two pyroxene thermometer (mean, mean - 1σ , and mean + 1σ), mineral modes, and REE analyses (IBM: 7 modes, 3 analyses; Tonga 6 modes, 3 analyses). All combinations were used to correct the data over 1000–1100 °C with 10 °C increments, resulting in 3528 corrected datasets for the IBM harzburgite and 2268 for the Tonga lherzolite sample. We then calculated the mean and standard deviations to obtain the representative data to compare with the modelled clinopyroxene (Fig. S-4). Average compositions of FAB from the Ogasawara area in the IBM region (Ishizuka *et al.*, 2011; Shervais *et al.*, 2019) and EAT from the Tonga region (Todd *et al.*, 2012) were calculated using samples with MgO contents greater than 7.5 wt.%, respectively (Fig. S-4).




Figure S-4 Chondrite normalised REE patterns of early basalts and subsolidus-corrected clinopyroxene from the studied samples (IBM harzburgite: ODP125-778A-003R-CC-4–7, Tonga lherzolite: BMRG08-111-3-6). **(a)** the IBM region and **(b)** the Tonga region. Symbols represent average values, and error bars indicate ± 2 standard deviations.

Influx-Free Open System Melting Models

We conducted a one dimensional, steady state open system melting model (Ozawa and Shimizu, 1995; Ozawa, 2001), where the full set of governing equations is described. The open system melting model encompasses batch melting, fractional melting, critical melting, melt–rock reactions, and fluid/melt flux melting, and thus it allows us to evaluate melt segregation and interaction with fluxed melt or fluids.

We conducted a suite of influx-free open system melting models to simulate REE compositions in accumulated melt and residual clinopyroxene. We used a depleted-MORB mantle (DMM) source or a depleted DMM (DDMM) source (Workman and Hart, 2005) (Table S-1). Melting mineral modes in the garnet peridotite stability field were from Walter (1998), and those in the spinel peridotite stability field were from

Kinzler and Grove (1992). Mineral modes from garnet peridotite to spinel peridotite were converted using the stoichiometry from Takazawa *et al.* (1996). We used mineral-melt partition coefficients from Kelemen *et al.* (2004), Warren (2016) and Sun and Liang (2014) (Table S-2). We first conducted influx-free open-system melting models to examine the link between our sample and early basalts (Table S-2), and we varied two key parameters (Table S-3): the critical melt fraction (a_c : 0–2%) at which melt starts to segregate, and the degree of melting (F : 0 up to 25% or until the residual clinopyroxene is exhausted). We varied the melting degree at the garnet peridotite stability field ($F_{grt} = 0$ –10%) prior to melting at the spinel peridotite stability field. We set $\gamma = 1$ for the melt separation rate during melting and a large τ value (= 10,000) for melt separation during trapped melt crystallisation, effectively neglecting the change of melt separation rate during melting and the stage of trapped melt crystallisation. We calculated the REE contents of clinopyroxene and the accumulated total melt at every 0.1% melting increment, resulting in over 20000 modelled clinopyroxene and melt compositions (Fig. S-5 and Fig. S-6).

Table S-1 Concentrations (ppm) of REE and mineral modes of starting material (DMM and DDMM sources) and melting modes for garnet peridotite stability field and spinel peridotite stability field.

Elements	DMM	DDMM
La	0.192	0.134
Ce	0.55	0.421
Pr	0.107	0.087
Nd	0.581	0.483
Sm	0.239	0.21
Eu	0.096	0.086
Gd	0.358	0.324
Tb	0.07	0.064
Dy	0.505	0.471
Ho	0.115	0.108
Er	0.348	0.329
Yb	0.365	0.348
Lu	0.058	0.056
Mineral Mode (%)	DMM	DDMM
Olivine	59	59.3
Orthopyroxene	22	22.1
Clinopyroxene	11	10.8
Garnet	8	7.8
Melting Mode	Garnet peridotite	
Olivine	0.04	
Orthopyroxene	-0.19	
Clinopyroxene	1.05	
Garnet	0.1	
Melting Mode	Spinel peridotite	
Olivine	-0.06	
Orthopyroxene	0.28	
Clinopyroxene	0.67	
Spinel	0.11	

Table S-2 Partition coefficients used for our melting models. We set the partition coefficients for spinel zero, following Warren (2016).

Element	Kd _{ol}	Kd _{cpx}	Kd _{opx}	Kd _{grt}	Kd _{amp}
La	0.0000023	0.0550	0.0007	0.001	0.112
Ce	0.0000073	0.0876	0.0016	0.008	0.189
Pr	0.000021	0.1318	0.0032	0.033	0.294
Nd	0.000058	0.1878	0.0060	0.057	0.42
Sm	0.00029	0.3083	0.0158	0.217	0.645
Eu	0.00055	0.3638	0.0227	0.45	0.72
Gd	0.0010	0.4169	0.0315	0.9	0.769
Tb	0.0017	0.4645	0.0422	1.5	0.786
Dy	0.0029	0.5034	0.0549	2	0.77
Ho	0.0045	0.5294	0.0680	2.8	0.729
Er	0.0066	0.5437	0.0808	3.5	0.673
Yb	0.0121	0.5453	0.1036	7	0.548
Lu	0.0153	0.5373	0.1128	9	0.491

Table S-3 A summary of key input parameters for influx-free open system melting models.

Model Setup	
Figures	Fig. 4 ($\beta = 0$) Fig. S-5, S-6, S-7, S-8
Source	DMM or DDMM
Critical melt fraction (a_c)	0–2%
Melting at the garnet peridotite stability field prior to melting at the spinel peridotite stability field (F_{grt})	0–10%
Total melting degree (F)	0 to 25%, or until the residual clinopyroxene (Cpx) is exhausted.
Melting degree step	0.1%
Clinopyroxene / Accumulate melt outputs	N = 21620

Our influx-free open-system melting models show that REE concentrations in clinopyroxene decrease with increasing F (Fig. S-5). At a given F , increasing a_c suppresses LREE depletion, whereas increasing F_{grt} suppresses HREE depletion in clinopyroxene from spinel peridotite, as partition coefficients of garnet for HREE are larger than 1. LREE depletion is thus strongly controlled by a_c , and HREE depletion is controlled by F_{grt} . This observation is consistent with the previous studies (Ozawa, 2001; Hellebrand *et al.*, 2002).




Figure S-5 Chondrite normalised REE patterns of modelled clinopyroxene from nine representative influx-free open system melting models from the DMM source. Each row corresponds to the melting degree in garnet peridotite stability field, F_{grt} , increasing from the top to bottom (0, 0.04, 0.1 = 0%, 4%, 10%). Each column represents a different critical melt fraction, a_c , increasing from left to right (0.001, 0.01, 0.02 = 0.1%, 1%, 2%). Colour indicates the degree of melting, F . Model results with low HREE contents correspond to clinopyroxene in the garnet peridotite stability field.

Accumulated melt is produced once melt separation begins, resulting in LREE enriched patterns at low a_c and low F (Fig. S-6). At a given F ($< 10\%$), accumulated melts generated in the garnet peridotite stability field are lower in HREE than melts generated in the spinel peridotite stability field. At given F ($> 10\%$), accumulated melt shows similar REE patterns regardless of a_c and F_{grt} , while those derived from the DDMM source exhibit slightly lower REE concentrations compared to those from the DMM source. Modal Ce contents in accumulated melts are 2.98 ppm for models derived from the DDMM source and 3.86 ppm for those derived from the DMM source (Fig. S-7). The difference in source Ce between DDMM (0.421 ppm) and DMM (0.55 ppm) is 0.13 ppm, which results in a ~ 0.88 ppm difference in the modal Ce contents in accumulated melts. Notably, the average Ce content in FAB (3.1 ppm) is closer to the mode of the DDMM-derived accumulated melts.




Figure S-6 Chondrite normalised REE patterns of modelled accumulated melt from nine representative influx-free open system melting models from the DMM source. The models presented here use the same parameters as in Figure S-5.




Figure S-7 Histogram of chondrite normalised Ce value (Ce_N) of modelled accumulated melt from all influx-free open system melting models, comparing models derived from the DMM and DDMM sources with FAB and EAT. Dotted lines indicate the modal Ce concentrations, estimated in log space, for models with DMM and DDMM sources. Symbols for the samples represent average values, and error bars indicate ± 2 standard deviations. The corresponding average Ce concentrations (ppm) are also indicated for clarity.

Best Fit Model Estimation

To evaluate how well the modelled REE contents match the REE contents in samples, we calculate the reduced chi-square (χ_v^2):

$$\chi_v^2 = \frac{1}{v} \cdot \sum \frac{(O_i - C_i)^2}{\sigma_i^2} \quad (\text{Eq. S-3})$$

χ_v^2 is the reduced chi-square, where v is the degrees of freedom, which is the number of elements minus the number of fitted parameters, O_i represents the mean concentration of element i in the sample, σ_i is the corresponding standard deviation, and C_i is the concentration of element i in the modelled data. χ_v^2 represents the similarity between two compositions in REE space, with a value close to 1 indicating a good fit, and we considered models with $\chi_v^2 < 1 + \sqrt{2/v}$ to represent good fits (Press *et al.*, 2007).

We first compared the modelled accumulated melts with the REE compositions of FAB and EAT, respectively, and estimated best fit models that reproduce the REE patterns of FAB and EAT based on the criteria defined above. Within each region, we subsequently performed a best fit estimation by comparing modelled clinopyroxene compositions with the corrected REE compositions of clinopyroxene, using the basalt based best fit models to evaluate whether the studied samples represent residues of the corresponding basaltic melts.

We obtained the best fit models reproducing both clinopyroxene and early basalt compositions (Fig. S-8). For the IBM harzburgite and FAB, the best fits correspond to $F = 16.8 \pm 2.1\%$ (2σ) with $a_c = 0.4 \pm 0.6\%$ and $F_{grt} = 2.9 \pm 3.8\%$. For the Tonga lherzolite and EAT, the best fits correspond to $F = 7.4\% \pm 0.1\%$ with $a_c = 1\%$ and $F_{grt} = 4\%$. Among IBM best fit models, 50.9% adopt the DDMM source and 40.9% the DMM source, whereas all Tonga best fit models are derived from the DMM source. Notably, all best fit models for the IBM clinopyroxene exhibit LREE depleted patterns, consistent with the samples in which LREE are below detection limits. The accumulated melts for the IBM FAB show slight differences in LREE contents, with models derived from the DDMM source yielding lower values than those from the DMM source.




Figure S-8 Chondrite normalised REE patterns of best-fit models compared with early basalts and subsolidus corrected clinopyroxene from the studied samples (IBM harzburgite: ODP125-778A-003R-CC-4–7, Tonga lherzolite: BMRG08-111-3-6). **(a)** the IBM region and **(b)** the Tonga region. Symbols for the samples represent average values, and error bars indicate ± 2 standard deviations. Mean and two standard deviations of the total melting degree (F), the critical melt fraction (a_c), and melting degree in the garnet peridotite stability field (F_{grt}) are indicated. Colour indicates the source type of each best-fit model (either DDMM or DMM).

Slab Derived Fluid Influx Open System Melting Models

Based on the best fit models estimated by influx-free open system melting models for each region, we conducted slab derived fluid influx open system melting models to investigate the effect of fluids and formation of amphibole (Table S-4). Slab derived fluid (or melt) compositions were adapted from Li *et al.* (2013) and Shervais *et al.* (2019), sourced from sediment and amphibolite (altered oceanic crust) (Table S-5). We set $a_c = 0.4\%$ and $F_{grt} = 4\%$ and varied the influx rate (β) from 0 to 1 for the IBM sample (Fig. S-9). We set $a_c = 1\%$ and $F_{grt} = 4\%$ and $\beta = 0$ to 1 for the Tonga sample (Fig. S-10). β is the mass fraction of influxed material relative to the initial solid over F . For example, when $F = 10\%$ and $\beta = 0.001$ or 1, influxed mass material is 0.01% and 10%, respectively.

To examine the formation of amphibole, we considered the crystallisation of trapped melt. We assume crystallisation of olivine, clinopyroxene, and amphibole in fixed proportions of 1:4:4. Partition coefficients for amphibole are taken from Shimizu *et al.* (2017) (Table S-2). We fixed the trapped melt fraction prior to crystallisation at 0.1% and the all trapped melt is assumed to crystallise in residual solid by setting $\tau = 0$ (*i.e.*, no melt escape during trapped melt crystallisation). The trapped melt fraction (0.1%) was therefore converted into solid phases (olivine, clinopyroxene, and amphibole) and equilibrated with the residual assemblage.

Using the parameters defined above, we modelled accumulated melt, residual clinopyroxene, and amphibole compositions up to $F = 25\%$, at increments of 0.1% melting degree.

Table S-4 A summary of key input parameters for slab derived fluid influx open system melting models.

Model Setup	For IBM	For Tonga
Figures	Fig. 4a–b, Fig. S-9, Fig. S-11	Fig. 4c–e, Fig. S-10, Fig. S-11
Source	DDMM	DMM
Critical melt fraction (a_c)	0.4%	1%
Melting at the garnet peridotite stability field prior to melting at the spinel peridotite stability field (F_{grt})		4%
Total melting degree (F)		0–25%
Influx composition	Slab derived fluid (Li <i>et al.</i> , 2013; Shervais <i>et al.</i> , 2019)	
Influx rate (β)	0 (influx-free), 0.0001, 0.001, 0.01, 0.1, 1	

Table S-5 REE contents (ppm) in the slab derived fluid used for our models.

Element	
La	9.3
Ce	27.4
Pr	2.97
Nd	11.3
Sm	1.7
Eu	0.46
Gd	1.17
Tb	0.16
Dy	0.91
Ho	0.177
Er	0.499
Yb	0.515
Lu	0.08

At the best fit F (IBM: 16.8%, Tonga: 7.4%), increasing β suppresses LREE depletion in clinopyroxene, resulting in flatter REE patterns and higher absolute LREE concentrations (Fig. S-9 and Fig. S-10). Amphibole exhibits a comparable response (Fig. S-10i–l). LREE contents in clinopyroxene and amphibole are highly sensitive to small variations in β within the range of 0–0.001, but their HREE contents decrease with increasing F (Fig. S-9e–g, S-10e–g, and S-10i–k). At β is high (>0.1), even HREE depletion in clinopyroxene and amphibole is suppressed (Fig. S-9h, S-10h, and S-10l). The effect of fluid influx on the accumulated melt is comparatively minor at low β (<0.01) (Fig. S-9 and S-10). At β is high ($=1$), the accumulated melt exhibits LREE enriched patterns, while HREE remain broadly similar to those in models with lower β .

At higher β (~ 0.01) and F ($>20\%$), the models produce LREE enriched yet HREE depleted patterns in clinopyroxene (Fig. S-9g and S-10g). These signatures resemble those observed in depleted forearc peridotites (Birner *et al.*, 2017; Loocke and Snow, 2024) and in ultra-depleted peridotites associated with boninitic magmatism, where slab derived fluid influx melting plays a key role (Nishio *et al.*, 2023).

Slab derived fluid influx open system melting models (IBM)

(a–d) Accumulated melt vs. FAB




Figure S-9 Chondrite normalised REE patterns of (a–d) the modelled accumulated melt and (e–h) modelled clinopyroxene from four representative slab derived fluid influx open system melting models from the DDMM source, comparing with FAB and subsolidus corrected clinopyroxene from the studied samples (IBM harzburgite: ODP125-778A-003R-CC-4–7). Each column represents a different influx rate, β , increasing from left to right (0, 0.0001, 0.01, 1). Colour indicates the degree of melting, F . Symbols for the samples represent average values, and error bars indicate ± 2 standard deviations.




Slab derived fluid influx open system melting models (Tonga)**(a–d) Accumulated melt vs. EAT****(e–h) Modelled Clinopyroxene vs. Sample Clinopyroxene****(i–l) Modelled Amphibole vs. Sample Amphibole**

Figure S-10 Chondrite normalised REE patterns of **(a–d)** the modelled accumulated melt and **(e–h)** modelled clinopyroxene, and **(i–l)** modelled amphibole from four representative slab derived fluid influx open system melting models from the DMM source, comparing with EAT, subsolidus corrected clinopyroxene, and amphibole from the studied samples (Tonga lherzolite: BMRG08-111-3-6). Each column represents a different influx rate, β , increasing from left to right (0, 0.0001, 0.01, 1). Colour indicates the degree of melting, F .

Figure S-11 presents REE patterns of best fit models of each region (IBM: $F = 16.8\%$, $a_c = 0.4\%$, $F_{grt} = 4\%$; Tonga: $F = 7.4\%$, $a_c = 1\%$, $F_{grt} = 4\%$), with varying β . At the best fit conditions, increasing β shows systematic changes in REE patterns. In both clinopyroxene and amphibole, higher β suppresses LREE depletion, whereas HREE remain relatively constant due to the low HREE concentrations in the slab derived fluid (Table S-5). Because the best fit F is higher for the IBM samples than for the Tonga samples, the suppression of LREE depletion is more pronounced at much lower β . For the IBM clinopyroxene, the detection limit of Ce corresponds to ~ 0.001 in chondrite normalised value, implying that β is likely < 0.0001 . In contrast, for the Tonga clinopyroxene, changes in REE patterns occur at relatively higher β (> 0.1).

Accumulated melt compositions at the best fit conditions are only weakly affected at low β (< 0.1), which indicates that clinopyroxene and amphibole are much more sensitive to small amounts of fluid influx.




Figure S-11 Chondrite normalised REE patterns: (a) modelled accumulated melt compared with FAB, (b) modelled clinopyroxene compared with subsolidus corrected IBM clinopyroxene, (c) modelled accumulated melt compared with EAT, (d) modelled clinopyroxene compared with subsolidus corrected Tonga clinopyroxene, and (e) modelled amphibole compared with Tonga amphibole. Sample symbols represent average values, and error bars indicate ± 2 standard deviations. All models are produced at fixed melting degree (F), critical melt fraction (a_c), and source for each region, as indicated at the top of each column. Model colour indicates the influx rate (β).

Compiled Forearc Spinel Data and Reference Data for Plots

We compiled spinel data from forearc peridotites (Table S-13), which are used in Figure 1. Data are from the IBM region (Bloomer and Hawkins, 1983; Ishii *et al.*, 1992; Parkinson and Pearce, 1998; Ishii *et al.*, 2000; D'Antonio and Kristensen, 2004; Okamura *et al.*, 2006; Zanetti *et al.*, 2006; Dare *et al.*, 2009; Murata *et al.*, 2009; Pabst, 2009; Morishita *et al.*, 2011; Harigane *et al.*, 2013; Ghosh, 2018; Debret *et al.*, 2019; Park, 2019; Albers *et al.*, 2020; Golich and Vysotskiy, 2020; Gose and Schmädicke, 2021; Ichiyama *et al.*, 2021; Miladinova *et al.*, 2023; Wang *et al.*, 2023) and the Tonga region (Birner *et al.*, 2016, 2017; Day and Brown, 2021). Spinel data from the South Mariana region were not included due to the tectonic complexity of these areas (Michibayashi *et al.*, 2009).

Reference clinopyroxene data used in Figure 2 are from the fertile residual abyssal peridotites, characterised by low spinel Cr# (0.10–0.11) and high LREE/HREE ratios in clinopyroxene, are from Johnson *et al.* (1990), Salters and Dick (2002), Seyler and Bonatti (1997), and Warren (2016). The LREE-depleted residual abyssal peridotites from Harigane *et al.* (2016). Clinopyroxene data of IBM peridotites are from Ichiyama *et al.* (2021), Ishii *et al.* (1992), Loocke and Snow (2024), and Parkinson and Pearce (1998). Clinopyroxene data of Tonga peridotites are from Birner *et al.* (2017) and Day and Brown (2021).

Reference amphibole data (Table S-14 and S-15) used for Figure 3 and S-12 are derived from IBM forearc peridotites (Ishii *et al.*, 1992; Parkinson and Pearce, 1998; Morishita *et al.*, 2011; Ichiyama *et al.*, 2021), ultra-depleted peridotites from the Kamuikotan unit, Japan (Nishio *et al.*, 2023; Tamura *et al.*, 2024), backarc peridotites (Akizawa *et al.*, 2021; Sen *et al.*, 2021; Gong *et al.*, 2022), abyssal peridotites (Seyler *et al.*, 2004; Cipriani *et al.*, 2009; Zhang *et al.*, 2024), and basal lherzolites from the Oman ophiolite (Prigent *et al.*, 2018). For comparison, we also show the compositional ranges of mid-ocean ridge basalt (MORB), backarc basin basalt (BABB), forearc basalt and early arc tholeiite (FAB-EAT), and boninite (Brounce *et al.*, 2021; Cottrell *et al.*, 2022).

The graphical abstract is adapted from Shuck *et al.* (2022) and modified to reflect the early basaltic magmatism stage.




Figure S-12 TiO_2 - K_2O discrimination plot for amphibole (Ozawa, 1988) from (a) forearc and ultra-depleted peridotites, (b) backarc peridotites, (c) abyssal peridotites, and (d) basal lherzolites from the Oman ophiolite. We plotted data with $\text{SiO}_2 < 52$ wt.%. Compositional area for mid-ocean ridge basalt (MORB), backarc basin basalt (BABB), forearc basalt and early arc tholeiite (FAB-EAT), and boninite are presented for comparison (Brounce *et al.*, 2021; Cottrell *et al.*, 2022). For amphibole from forearc peridotites, data from the IBM (Ogasawara), IBM (Conical), and IBM (Torishima) regions are from Morishita *et al.* (2011), Parkinson and Pearce (1998), and Ishii *et al.* (1992), respectively. Data for ultra-depleted peridotites from the Kamuikotan unit are from Nishio *et al.* (2023) and Tamura *et al.* (2024). For backarc peridotites, data from the Parece Vela Basin and Shikoku Basin are from Gong *et al.* (2022), and Akizawa *et al.* (2021) and Sen *et al.* (2021), respectively. For abyssal peridotites, SWIR plagioclase lherzolite data are from Zhang *et al.* (2024), SWIR harzburgite data are from Seyler *et al.* (2004), and MAR peridotite data are from Cipriani *et al.* (2009). Data for basal lherzolites from the Oman ophiolite are from Prigent *et al.* (2018). Amphibole occurrences in IBM (Ogasawara) in Spl and Kamuikotan (ultra-depleted) in Spl peridotites represent inclusions hosted by spinel. The detection limit for K_2O in amphibole in our analyses is ~ 0.1 wt.%.

Supplementary Tables

Table S-6	Sample information.
Table S-7	Mineral major element chemistry.
Table S-8	Mineral trace elements contents.
Table S-9	Corrected and/or representative REE concentrations in samples.
Table S-10	Results of representative influx-free open system melting models.
Table S-11	Best fit influx-free open system melting models.
Table S-12	Result of slab derived fluid influx open system melting models.
Table S-13	Compiled spinel data from forearc peridotites.
Table S-14	Compiled amphibole major element data.
Table S-15	Compiled amphibole REE data.

Tables S-6 to S-15 (.xlsx) are available for download from the online version of this article at <https://doi.org/10.7185/geochemlet.2624>.

Supplementary Information References

- Akizawa, N., Ozawa, K., Tamura, A., Michibayashi, K., Arai, S. (2016) Three-dimensional Evolution of Melting, Heat and Melt Transfer in Ascending Mantle beneath a Fast-spreading Ridge Segment Constrained by Trace Elements in Clinopyroxene from Concordant Dunites and Host Harzburgites of the Oman Ophiolite. *Journal of Petrology* 57, 777–814. <https://doi.org/10.1093/petrology/egw020>
- Akizawa, N., Ohara, Y., Okino, K., Ishizuka, O., Yamashita, H., Machida, S., Sanfilippo, A., Basch, V., Snow, J.E., Sen, A., Hirauchi, K.-i., Michibayashi, K., Harigane, Y., Fujii, M., Asanuma, H., Hirata, T. (2021) Geochemical characteristics of back-arc basin lower crust and upper mantle at final spreading stage of Shikoku Basin: an example of Mado Megamullion. *Progress in Earth and Planetary Science* 8, 65. <https://doi.org/10.1186/s40645-021-00454-3>
- Albers, E., Kahl, W.-A., Beyer, L., Bach, W. (2020) Variant across-forearc compositions of slab-fluids recorded by serpentinites: Implications on the mobilization of FMEs from an active subduction zone (Mariana forearc). *Lithos* 364–365, 105525. <https://doi.org/10.1016/j.lithos.2020.105525>
- Arai, S. (1984) Igneous mineral equilibria in some alpine-type peridotites in Japan. In: Sunagawa, I. (Ed.) *Materials Science of the Earth's Interior*. Springer, Dordrecht, 445–460.
- Birner, S.K., Warren, J.M., Cottrell, E., Davis, F.A. (2016) Hydrothermal alteration of seafloor peridotites does not influence oxygen fugacity recorded by spinel oxybarometry. *Geology* 44, 535–538. <https://doi.org/10.1130/G38113.1>
- Birner, S.K., Warren, J.M., Cottrell, E., Davis, F.A., Kelley, K.A., Falloon, T.J. (2017) Forearc Peridotites from Tonga Record Heterogeneous Oxidation of the Mantle following Subduction Initiation. *Journal of Petrology* 58, 1755–1780. <https://doi.org/10.1093/petrology/egx072>
- Bloomer, S.H., Hawkins, J.W. (1983) Gabbroic and Ultramafic Rocks from the Mariana Trench: An Island Arc Ophiolite. In: Hayes, D.E. (Ed.) *The Tectonic and Geologic Evolution of Southeast Asian Seas and Islands: Part 2*. American Geophysical Union, Washington, D.C., 294–317. <https://doi.org/10.1029/GM027p0294>

- Bloomer, S.H., Wright, D., MacLeod, C.J., Tappin, D., Clift, P., Fallon, T., Fisher, R.L., Gillis, K., Ishii, T., Kelman, M., Mafi, K., Sato, H., Winowitch, M. (1996) Geology of the Tonga Forearc: A supra-subduction zone ophiolite. *EOS, Transactions, American Geophysical Union Fall Meeting*, San Francisco, California, F325 [abstract].
- Brey, G.P., Köhler, T. (1990) Geothermobarometry in Four-phase Lherzolites II. New Thermobarometers, and Practical Assessment of Existing Thermobarometers. *Journal of Petrology* 31, 1353–1378. <https://doi.org/10.1093/petrology/31.6.1353>
- Brounce, M., Reagan, M.K., Kelley, K.A., Cottrell, E., Shimizu, K., Almeev, R. (2021) Covariation of Slab Tracers, Volatiles, and Oxidation During Subduction Initiation. *Geochemistry, Geophysics, Geosystems* 22, e2021GC009823. <https://doi.org/10.1029/2021GC009823>
- Cipriani, A., Bonatti, E., Seyler, M., Brueckner, H.K., Brunelli, D., Dallai, L., Hemming, S.R., Ligi, M., Ottolini, L., Turrin, B.D. (2009) A 19 to 17 Ma amagmatic extension event at the Mid-Atlantic Ridge: Ultramafic mylonites from the Vema Lithospheric Section. *Geochemistry, Geophysics, Geosystems* 10, Q10011. <https://doi.org/10.1029/2009GC002534>
- Cottrell, E., Birner, S.K., Brounce, M., Davis, F.A., Waters, L.E., Kelley, K.A. (2022) Oxygen Fugacity Across Tectonic Settings. In: Moretti, R., Neuville, D.R. (Eds.) *Magma Redox Geochemistry*. American Geophysical Union, Washington, D.C., 33–61. <https://doi.org/10.1002/9781119473206.ch3>
- D'Antonio, M., Kristensen, M.B. (2004) Serpentine and brucite of ultramafic clasts from the South Chamorro Seamount (Ocean Drilling Program Leg 195, Site 1200): inferences for the serpentinization of the Mariana forearc mantle. *Mineralogical Magazine* 68, 887–904. <https://doi.org/10.1180/0026461046860229>
- Dare, S.A.S., Pearce, J.A., McDonald, I., Styles, M.T. (2009) Tectonic discrimination of peridotites using fO_2 –Cr# and Ga–Ti–Fe^{III} systematics in chrome–spinel. *Chemical Geology* 261, 199–216. <https://doi.org/10.1016/j.chemgeo.2008.08.002>
- Day, J.M.D., Brown, D.B. (2021) Ancient Melt-Depletion in Fresh to Strongly Serpentinized Tonga Trench Peridotites. *Journal of Petrology* 62, egab088. <https://doi.org/10.1093/petrology/egab088>
- Debret, B., Albers, E., Walter, B., Price, R., Barnes, J.D., Beunon, H., Facq, S., Gillikin, D.P., Mattielli, N., Williams, H. (2019) Shallow forearc mantle dynamics and geochemistry: New insights from IODP Expedition 366. *Lithos* 326–327, 230–245. <https://doi.org/10.1016/j.lithos.2018.10.038>
- Dygert, N., Liang, Y. (2015) Temperatures and cooling rates recorded in REE in coexisting pyroxenes in ophiolitic and abyssal peridotites. *Earth and Planetary Science Letters* 420, 151–161. <https://doi.org/10.1016/j.epsl.2015.02.042>
- Ghosh, T. (2018) *Mantle Wedge Evolution Beneath the Northern Mariana Forearc*. Ph.D. Thesis, University of Houston.
- Golich, A.N., Vysotskiy, S.V. (2020) Island-Arc Ophiolites of the Hahajima Seamount (Bonin Trench, Philippine Sea). *Russian Journal of Pacific Geology* 14, 221–240. <https://doi.org/10.1134/S1819714020030033>
- Gong, X., Tian, L., Dong, Y. (2022) Contrasting melt percolation and melt-rock reactions in the Parece Vela back-arc oceanic lithosphere, Philippine Sea: A mineralogical perspective. *Lithos* 422–423, 106727. <https://doi.org/10.1016/j.lithos.2022.106727>
- Gose, J., Schmädicke, E. (2021) Water in the Supra-Subduction-Zone Mantle of the Mariana-Izu-Bonin Forearc: Constraints From Peridotitic Orthopyroxene. *Geochemistry, Geophysics, Geosystems* 22, e2020GC009586. <https://doi.org/10.1029/2020GC009586>
- Harigane, Y., Michibayashi, K., Morishita, T., Tani, K., Dick, H.J.B., Ishizuka, O. (2013) The earliest mantle fabrics formed during subduction zone infancy. *Earth and Planetary Science Letters* 377–378, 106–113. <https://doi.org/10.1016/j.epsl.2013.06.031>
- Harigane, Y., Abe, N., Michibayashi, K., Kimura, J.-I., Chang, Q. (2016) Melt-rock interactions and fabric development of peridotites from North Pond in the Kane area, Mid-Atlantic Ridge: Implications of microstructural and petrological analyses of peridotite samples from IODP Hole U1382A. *Geochemistry, Geophysics, Geosystems* 17, 2298–2322. <https://doi.org/10.1002/2016GC006429>
- Hellebrand, E., Snow, J.E., Hoppe, P., Hofmann, A.W. (2002) Garnet-field Melting and Late-stage Refertilization in ‘Residual’ Abyssal Peridotites from the Central Indian Ridge. *Journal of Petrology* 43, 2305–2338. <https://doi.org/10.1093/petrology/43.12.2305>
- Ichiyama, Y., Tsujimori, T., Fryer, P., Michibayashi, K., Tamura, A., Morishita, T. (2021) Temporal and spatial mineralogical changes in clasts from Mariana serpentinite mud volcanoes: Cooling of the hot forearc-mantle at subduction initiation. *Lithos* 384–385, 105941. <https://doi.org/10.1016/j.lithos.2020.105941>
- Ishii, T., Robinson, P.T., Maekawa, H., Fiske, R. (1992) Petrological studies of peridotites from diapiric serpentinite seamounts in the Izu-Ogasawara-Mariana Forearc, Leg 125. In: Fryer, P., Pearce, J.A., Stokking, L.B., Ali, J.R.,

- Arculus, R., *et al.* (Eds.) *Proceedings of the Ocean Drilling Program, Scientific Results*, 125, Ocean Drilling Program, College Station, TX, 445–485. <https://doi.org/10.2973/odp.proc.sr.125.129.1992>
- Ishii, T., Sato, H., Haraguchi, S., Fryer, P., Fujioka, K., Bloomer, S., Yokose, H. (2000) Petrological Characteristics of Peridotites from Serpentine Seamounts in the Izu-Ogasawara-Mariana Forearc. *Journal of Geography (Chigaku Zasshi)* 109, 517–530. https://doi.org/10.5026/jgeography.109.4_517
- Ishizuka, O., Tani, K., Reagan, M.K., Kanayama, K., Umino, S., Harigane, Y., Sakamoto, I., Miyajima, Y., Yuasa, M., Dunkley, D.J. (2011) The timescales of subduction initiation and subsequent evolution of an oceanic island arc. *Earth and Planetary Science Letters* 306, 229–240. <https://doi.org/10.1016/j.epsl.2011.04.006>
- Jochum, K.P., Weis, U., Stoll, B., Kuzmin, D., Yang, Q., Raczek, I., Jacob, D.E., Stracke, A., Birbaum, K., Frick, D.A., Günther, D., Enzweiler, J. (2011) Determination of Reference Values for NIST SRM 610–617 Glasses Following ISO Guidelines. *Geostandards and Geoanalytical Research* 35, 397–429. <https://doi.org/10.1111/j.1751-908X.2011.00120.x>
- Johnson, K.T.M., Dick, H.J.B., Shimizu, N. (1990) Melting in the oceanic upper mantle: An ion microprobe study of diopsides in abyssal peridotites. *Journal of Geophysical Research: Solid Earth* 95, 2661–2678. <https://doi.org/10.1029/JB095iB03p02661>
- Kelemen, P.B., Yogodzinski, G.M., Scholl, D.W. (2004) Along-Strike Variation in the Aleutian Island Arc: Genesis of High Mg# Andesite and Implications for Continental Crust. In: Eiler, J. (Ed.) *Inside the Subduction Factory*. American Geophysical Union, Washington, D.C., 223–276. <https://doi.org/10.1029/138GM11>
- Kinzler, R.J., Grove, T.L. (1992) Primary magmas of mid-ocean ridge basalts 1. Experiments and methods. *Journal of Geophysical Research: Solid Earth* 97, 6885–6906. <https://doi.org/10.1029/91JB02840>
- Lanari, P., Tedeschi, M. (2025) Chemical map classification in XMapTools. *Applied Computing and Geosciences* 25, 100230. <https://doi.org/10.1016/j.acags.2025.100230>
- Lanari, P., Vidal, O., De Andrade, V., Dubacq, B., Lewin, E., Grosch, E.G., Schwartz, S. (2014) XMapTools: A MATLAB®-based program for electron microprobe X-ray image processing and geothermobarometry. *Computers & Geosciences* 62, 227–240. <https://doi.org/10.1016/j.cageo.2013.08.010>
- Li, Y.-B., Kimura, J.-I., Machida, S., Ishii, T., Ishiwatari, A., Maruyama, S., Qiu, H.-N., Ishikawa, T., Kato, Y., Haraguchi, S., Takahata, N., Hirahara, Y., Miyazaki, T. (2013) High-Mg Adakite and Low-Ca Boninite from a Bonin Fore-arc Seamount: Implications for the Reaction between Slab Melts and Depleted Mantle. *Journal of Petrology* 54, 1149–1175. <https://doi.org/10.1093/petrology/egt008>
- Liang, Y., Sun, C., Yao, L. (2013) A REE-in-two-pyroxene thermometer for mafic and ultramafic rocks. *Geochimica et Cosmochimica Acta* 102, 246–260. <https://doi.org/10.1016/j.gca.2012.10.035>
- Liang, Y., Ji, Z., Liu, B. (2021) What can we learn from REE abundances in clinopyroxene and orthopyroxene in residual mantle peridotites? *Contributions to Mineralogy and Petrology* 176, 24. <https://doi.org/10.1007/s00410-021-01780-x>
- Lin, K.-Y., Warren, J.M., Davis, F.A. (2023) Trace elements in abyssal peridotite olivine record melting, thermal evolution, and melt refertilization in the oceanic upper mantle. *Contributions to Mineralogy and Petrology* 178, 66. <https://doi.org/10.1007/s00410-023-02044-6>
- Longerich, H.P., Jackson, S.E., Günther, D. (1996) Inter-laboratory note. Laser ablation inductively coupled plasma mass spectrometric transient signal data acquisition and analyte concentration calculation. *Journal of Analytical Atomic Spectrometry* 11, 899–904. <https://doi.org/10.1039/JA9961100899>
- Loocke, M.P., Snow, J.E. (2024) Arc Foundations and Subduction Initiation: Insights into the Magmatic Evolution of the Lower Crust/Upper Mantle of the Izu–Bonin Forearc during Subduction Initiation. *Journal of Petrology* 65, egae038. <https://doi.org/10.1093/petrology/egae038>
- Michibayashi, K., Ohara, Y., Stern, R.J., Fryer, P., Kimura, J.-I., Tasaka, M., Harigane, Y., Ishii, T. (2009) Peridotites from a ductile shear zone within back-arc lithospheric mantle, southern Mariana Trench: Results of a *Shinkai* 6500 dive. *Geochemistry, Geophysics, Geosystems* 10, Q05X06. <https://doi.org/10.1029/2008GC002197>
- Miladinova, I., Kurz, W., Hilmbauer-Hofmarcher, T. (2023) Fluid-Mantle Interaction Along the Mariana Convergent Margin. *Geochemistry, Geophysics, Geosystems* 24, e2023GC010968. <https://doi.org/10.1029/2023GC010968>
- Morishita, T., Tani, K., Shukuno, H., Harigane, Y., Tamura, A., Kumagai, H., Hellebrand, E. (2011) Diversity of melt conduits in the Izu-Bonin-Mariana forearc mantle: Implications for the earliest stage of arc magmatism. *Geology* 39, 411–414. <https://doi.org/10.1130/G31706.1>
- Murata, K., Maekawa, H., Yokose, H., Yamamoto, K., Fujioka, K., Ishii, T., Chiba, H., Wada, Y. (2009) Significance of serpentinization of wedge mantle peridotites beneath Mariana forearc, western Pacific. *Geosphere* 5, 90–104. <https://doi.org/10.1130/GES00213.1>
- Nishio, I., Morishita, T., Tamura, A., Itano, K., Takamizawa, S., Ichiyama, Y., Arai, S., Barrett, N., Szilas, K. (2023) Formation of Ultra-Depleted Mantle Peridotites and Their Relationship With Boninitic Melts: An Example

- From the Kamuikotan Unit, Hokkaido, Japan. *Journal of Geophysical Research: Solid Earth* 128, e2022JB025066. <https://doi.org/10.1029/2022JB025066>
- Okamura, H., Arai, S., Kim, Y.-U. (2006) Petrology of forearc peridotite from the Hahajima Seamount, the Izu-Bonin arc, with special reference to chemical characteristics of chromian spinel. *Mineralogical Magazine* 70, 15–26. <https://doi.org/10.1180/0026461067010310>
- Ozawa, K. (1988) Ultramafic tectonite of the Miyamori ophiolitic complex in the Kitakami Mountains, Northeast Japan: hydrous upper mantle in an island arc. *Contributions to Mineralogy and Petrology* 99, 159–175. <https://doi.org/10.1007/BF00371458>
- Ozawa, K. (2001) Mass balance equations for open magmatic systems: Trace element behavior and its application to open system melting in the upper mantle. *Journal of Geophysical Research: Solid Earth* 106, 13407–13434. <https://doi.org/10.1029/2001JB900001>
- Ozawa, K., Shimizu, N. (1995) Open-system melting in the upper mantle: Constraints from the Hayachine-Miyamori ophiolite, northeastern Japan. *Journal of Geophysical Research: Solid Earth* 100, 22315–22335. <https://doi.org/10.1029/95JB01967>
- Pabst, S. (2009) *Investigation of blueschist and serpentized harzburgite from the Mariana forearc: Insights into the mechanisms of element mobilization in subduction zones and storage of fluid-mobile elements in the mantle wedge*. Ph.D. Thesis, Heidelberg University.
- Park, J.-h. (2019) *Influence of partial melting and melt-mantle interaction on PGE geochemistry of the Mariana forearc peridotites*. Ph.D. Thesis, Seoul National University.
- Parkinson, I.J., Pearce, J.A. (1998) Peridotites from the Izu–Bonin–Mariana Forearc (ODP Leg 125): Evidence for Mantle Melting and Melt–Mantle Interaction in a Supra-Subduction Zone Setting. *Journal of Petrology* 39, 1577–1618. <https://doi.org/10.1093/petroj/39.9.1577>
- Parkinson, I.J., Pearce, J.A., Thirlwall, M.F., Johnson, K.T.M., Ingram, G. (1992) Trace element geochemistry of peridotites from the Izu-Bonin-Mariana Forearc, Leg 125. In: Fryer, P., Pearce, J.A., Stokking, L.B., Ali, J.R., Arculus, R., *et al.* (Eds.) *Proceedings of the Ocean Drilling Program, Scientific Results*, 125, Ocean Drilling Program, College Station, TX, 487–506. <https://doi.org/10.2973/odp.proc.sr.125.183.1992>
- Pearce, N.J.G., Perkins, W.T., Westgate, J.A., Gorton, M.P., Jackson, S.E., Neal, C.R., Chenery, S.P. (1997) A Compilation of New and Published Major and Trace Element Data for NIST SRM 610 and NIST SRM 612 Glass Reference Materials. *Geostandards Newsletter* 21, 115–144. <https://doi.org/10.1111/j.1751-908X.1997.tb00538.x>
- Poldervaart, A., Hess, H.H. (1951) Pyroxenes in the Crystallization of Basaltic Magma. *The Journal of Geology* 59, 472–489. <https://doi.org/10.1086/625891>
- Press, W.H., Teukolsky, S.A., Vetterling, W.T., Flannery, B.P. (2007) *Numerical Recipes: The Art of Scientific Computing*. Third Edition, Cambridge University Press, Cambridge.
- Prigent, C., Agard, P., Guillot, S., Godard, M., Dubacq, B. (2018) Mantle Wedge (De)formation During Subduction Infancy: Evidence from the Base of the Semail Ophiolitic Mantle. *Journal of Petrology* 59, 2061–2092. <https://doi.org/10.1093/petrology/egy090>
- Sagawa, T., Tamura, A., Okino, K., Morishita, T. (2022) Validation of Trace Element Analysis of Geological Materials by Single-Pulse Laser Ablation-Inductively Coupled Plasma-Mass Spectrometry (LA-ICP-MS). *Analytical Letters* 55, 2886–2896. <https://doi.org/10.1080/00032719.2022.2077356>
- Salter, V.J.M., Dick, H.J.B. (2002) Mineralogy of the mid-ocean-ridge basalt source from neodymium isotopic composition of abyssal peridotites. *Nature* 418, 68–72. <https://doi.org/10.1038/nature00798>
- Sen, A., Snow, J.E., Ohara, Y., Hirauchi, K., Kouketsu, Y., Sanfilippo, A., Basch, V., Harigane, Y., Fujii, M., Okino, K., Akizawa, N. (2021) Melting and Evolution of Amphibole-Rich Back-Arc Abyssal Peridotites at the Mado Megamullion, Shikoku Basin. *Geochemistry, Geophysics, Geosystems* 22, e2021GC010013. <https://doi.org/10.1029/2021GC010013>
- Seyler, M., Bonatti, E. (1997) Regional-scale melt-rock interaction in lherzolitic mantle in the Romanche Fracture Zone (Atlantic Ocean). *Earth and Planetary Science Letters* 146, 273–287. [https://doi.org/10.1016/S0012-821X\(96\)00220-8](https://doi.org/10.1016/S0012-821X(96)00220-8)
- Seyler, M., Lorand, J.-P., Toplis, M.J., Godard, G. (2004) Asthenospheric metasomatism beneath the mid-ocean ridge: Evidence from depleted abyssal peridotites. *Geology* 32, 301–304. <https://doi.org/10.1130/G20191.1>
- Shervais, J.W., Reagan, M., Haugen, E., Almeev, R.R., Pearce, J.A., Prytulak, J., Ryan, J.G., Whattam, S.A., Godard, M., Chapman, T., Li, H., Kurz, W., Nelson, W.R., Heaton, D., Kirchenbaur, M., Shimizu, K., Sakuyama, T., Li, Y., Vetter, S.K. (2019) Magmatic Response to Subduction Initiation: Part 1. Fore-arc Basalts of the Izu-Bonin Arc From IODP Expedition 352. *Geochemistry, Geophysics, Geosystems* 20, 314–338. <https://doi.org/10.1029/2018GC007731>

- Shimizu, K., Liang, Y., Sun, C., Jackson, C.R.M., Saal, A.E. (2017) Parameterized lattice strain models for REE partitioning between amphibole and silicate melt. *American Mineralogist* 102, 2254–2267. <https://doi.org/10.2138/am-2017-6110>
- Shuck, B., Gulick, S.P.S., Van Avendonk, H.J.A., Gurnis, M., Sutherland, R., Stock, J., Hightower, E. (2022) Stress transition from horizontal to vertical forces during subduction initiation. *Nature Geoscience* 15, 149–155. <https://doi.org/10.1038/s41561-021-00880-4>
- Sun, C., Liang, Y. (2012) Distribution of REE between clinopyroxene and basaltic melt along a mantle adiabat: effects of major element composition, water, and temperature. *Contributions to Mineralogy and Petrology* 163, 807–823. <https://doi.org/10.1007/s00410-011-0700-x>
- Sun, C., Liang, Y. (2013) The importance of crystal chemistry on REE partitioning between mantle minerals (garnet, clinopyroxene, orthopyroxene, and olivine) and basaltic melts. *Chemical Geology* 358, 23–36. <https://doi.org/10.1016/j.chemgeo.2013.08.045>
- Sun, C., Liang, Y. (2014) An assessment of subsolidus re-equilibration on REE distribution among mantle minerals olivine, orthopyroxene, clinopyroxene, and garnet in peridotites. *Chemical Geology* 372, 80–91. <https://doi.org/10.1016/j.chemgeo.2014.02.014>
- Takazawa, E., Frey, F., Shimizu, N., Obata, M. (1996) Evolution of the Horoman Peridotite (Hokkaido, Japan): Implications from pyroxene compositions. *Chemical Geology* 134, 3–26. [https://doi.org/10.1016/S0009-2541\(96\)00083-6](https://doi.org/10.1016/S0009-2541(96)00083-6)
- Tamura, A., Sagawa, T., Okino, K., Morishita, T. (2022) Determination of whole-rock trace-element compositions of siliceous rocks using MgO-diluted fused glass and LA-ICP-MS. *Geochemical Journal* 56, 231–239. <https://doi.org/10.2343/geochemj.GJ22020>
- Tamura, A., Nishio, I., Iwata, Y., Arai, S., Morishita, T. (2024) Amphibole in ultra-depleted peridotites from the Horokanai ophiolite, Hokkaido, Japan: Boninite melt trapped in the mantle wedge. *Ophioliti* 49, 37–54. <https://doi.org/10.4454/ofioliti.v49i1.568>
- Todd, E., Gill, J.B., Pearce, J.A. (2012) A variably enriched mantle wedge and contrasting melt types during arc stages following subduction initiation in Fiji and Tonga, southwest Pacific. *Earth and Planetary Science Letters* 335–336, 180–194. <https://doi.org/10.1016/j.epsl.2012.05.006>
- Walter, M.J. (1998) Melting of Garnet Peridotite and the Origin of Komatiite and Depleted Lithosphere. *Journal of Petrology* 39, 29–60. <https://doi.org/10.1093/ptro/39.1.29>
- Wang, Y., Deng, J., Liao, R., Chen, L., Li, D., Liu, H., Sun, W. (2023) Magnesium isotopic composition of the Mariana forearc serpentinite: Implications for Mg isotopic composition of the mantle wedge and Mg isotopic fractionation during mantle wedge serpentinization. *Chemical Geology* 624, 121428. <https://doi.org/10.1016/j.chemgeo.2023.121428>
- Warren, J.M. (2016) Global variations in abyssal peridotite compositions. *Lithos* 248–251, 193–219. <https://doi.org/10.1016/j.lithos.2015.12.023>
- Witt-Eickchen, G., O'Neill, H.St.C. (2005) The effect of temperature on the equilibrium distribution of trace elements between clinopyroxene, orthopyroxene, olivine and spinel in upper mantle peridotite. *Chemical Geology* 221, 65–101. <https://doi.org/10.1016/j.chemgeo.2005.04.005>
- Workman, R.K., Hart, S.R. (2005) Major and trace element composition of the depleted MORB mantle (DMM). *Earth and Planetary Science Letters* 231, 53–72. <https://doi.org/10.1016/j.epsl.2004.12.005>
- Wright, D.J., Bloomer, S.H., MacLeod, C.J., Taylor, B., Goodlife, A.M. (2000) Bathymetry of the Tonga Trench and Forearc: a map series. *Marine Geophysical Researches* 21, 489–512. <https://doi.org/10.1023/A:1026514914220>
- Yao, L., Sun, C., Liang, Y. (2012) A parameterized model for REE distribution between low-Ca pyroxene and basaltic melts with applications to REE partitioning in low-Ca pyroxene along a mantle adiabat and during pyroxenite-derived melt and peridotite interaction. *Contributions to Mineralogy and Petrology* 164, 261–280. <https://doi.org/10.1007/s00410-012-0737-5>
- Zanetti, A., D'Antonio, M., Spadea, P., Raffone, N., Vannucci, R., Bruguier, O. (2006) Petrogenesis of Mantle Peridotites from the Izu-Bonin-Mariana (IBM) Forearc. *Ophioliti* 31, 189–206. <https://doi.org/10.4454/ofioliti.v31i2.340>
- Zhang, W.-Q., Liu, C.-Z., Dick, H.J.B., Mitchell, R.N., Liu, B.-D. (2024) Oceanic mantle beneath ultraslow spreading ridges metasomatized by variably evolved melts. *Contributions to Mineralogy and Petrology* 179, 15. <https://doi.org/10.1007/s00410-023-02093-x>